

GABA_A Receptor Trafficking Is Regulated by Protein Kinase C ϵ and the *N*-Ethylmaleimide-Sensitive Factor

Wen-Hai Chou,¹ Dan Wang,¹ Thomas McMahon,¹ Zhan-Heng Qi,¹ Maengseok Song,¹ Chao Zhang,² Kevan M. Shokat,² and Robert O. Messing¹

¹Ernest Gallo Clinic and Research Center, Department of Neurology, University of California, San Francisco, Emeryville, California 94688, and ²Howard Hughes Medical Institute and Department of Cellular and Molecular Pharmacology, University of California, San Francisco, San Francisco, California 94143

Disturbances in GABA_A receptor trafficking contribute to several neurological and psychiatric disorders by altering inhibitory neurotransmission. Identifying mechanisms that regulate GABA_A receptor trafficking could lead to better understanding of disease pathogenesis and treatment. Here, we show that protein kinase C ϵ (PKC ϵ) regulates the *N*-ethylmaleimide-sensitive factor (NSF), an ATPase critical for membrane fusion events, and thereby promotes the trafficking of GABA_A receptors. Activation of PKC ϵ decreased cell surface expression of GABA_A receptors and attenuated GABA_A currents. Activated PKC ϵ associated with NSF, phosphorylated NSF at serine 460 and threonine 461, and increased NSF ATPase activity, which was required for GABA_A receptor downregulation. These findings identify new roles for NSF and PKC ϵ in regulating synaptic inhibition through downregulation of GABA_A receptors. Reducing NSF activity by inhibiting PKC ϵ could help restore synaptic inhibition in disease states in which it is impaired.

Introduction

GABA_A receptors are heteropentameric ligand-gated ion channels that mediate most fast inhibitory neurotransmission (Olsen and Sieghart, 2008). Impaired GABA_A receptor signaling contributes to several neuropsychiatric conditions including epilepsy (Goodkin et al., 2008), schizophrenia (Charych et al., 2009), anxiety and depression (Kalueff and Nutt, 2007; Charych et al., 2009), substance abuse (Kumar et al., 2004), and pain (Knabl et al., 2008). GABA_A receptors have therefore been important targets for drug development, although current drugs can produce limiting side effects including sedation, amnesia, and dependence. Understanding mechanisms that regulate GABA_A receptor trafficking could increase knowledge about the pathophysiology of these disease states and provide clues for developing alternative drugs that modulate inhibitory neurotransmission with fewer side effects.

Tumor-promoting phorbol esters decrease cell surface GABA_A receptors in several preparations (Leidenheimer et al., 1992; Leidenheimer and Chapell, 1997; Chapell et al., 1998; Connolly et al., 1999; Cinar and Barnes, 2001; Balduzzi et al., 2002; Meier et al., 2003; Herring et al., 2005). This effect is presumed to be caused by activation of protein kinase C (PKC)

(Kittler and Moss, 2003; Leidenheimer, 2008). PKC is not a single entity but a family of nine kinases with several splice variants that transduce signals involving lipid second messengers (Song and Messing, 2005). Phorbol esters activate seven isozymes (α , β , γ , δ , ϵ , η , and θ), each with specialized functions that are generally not redundant. In addition to PKCs, phorbol esters activate several other proteins such as protein kinase D, chimerins, RasGRP, and Munc-13 (Brose and Rosenmund, 2002). Thus, it is not certain that phorbol esters regulate GABA_A receptor trafficking by activating a specific PKC isozyme or another phorbol ester-responsive protein.

PKC substrates that mediate GABA_A receptor downregulation are also unknown. Although several phosphorylation sites have been identified on intracellular loops of GABA_A γ 2 and β subunits, alanine substitutions do not prevent receptor downregulation by phorbol esters (Chapell et al., 1998; Connolly et al., 1999). Other possible substrates include proteins associated with GABA_A receptors (Chen and Olsen, 2007). One candidate is the *N*-ethylmaleimide-sensitive factor (NSF) (Matveeva et al., 2001), an ATPase that regulates membrane fusion events (Zhao et al., 2007) and is important for stability of AMPA receptors at excitatory synapses (Hanley et al., 2002). When overexpressed, NSF can decrease cell surface levels of GABA_A receptors (Goto et al., 2005). NSF also interacts with GABARAP (GABA_A receptor-associated protein), and this interaction is proposed to modulate GABA_A receptor trafficking (Kittler et al., 2001; Leil et al., 2004; Marsden et al., 2007). However, the hypothesis that endogenous NSF truly regulates GABA_A receptor trafficking remains to be tested.

We found that mice lacking PKC ϵ (*Prkce*^{-/-} mice) are hypersensitive to drugs that increase GABA_A receptor activation, including benzodiazepines, ethanol, neurosteroids, and barbiturates (Hodge et al., 1999, 2002). Because diminished

Received Jan. 17, 2010; revised Aug. 18, 2010; accepted Aug. 24, 2010.

This work was supported by United States Public Health Service Grant AA013588 and funds provided by the State of California for medical research on alcohol and substance abuse through University of California, San Francisco (UCSF) (R.O.M.). We thank S. Whiteheart (University of Kentucky) for the NSF cDNA construct, P. Sung and P. Chi (Yale University) for advice on the ATPase activity assay, Y. Li (University of Texas Southwestern) for mass spectrometry, C. Johnston (UCSF) for statistical advice, and A. Newton (University of California, San Diego) for the MyrPalm-CFP expression plasmid.

Correspondence should be addressed to Dr. Robert O. Messing, Ernest Gallo Clinic and Research Center, 5858 Horton Street, Suite 200, Emeryville, CA 94688. E-mail: romes@gallo.ucsf.edu.

W.-H. Chou's present address: Department of Biological Sciences, Kent State University, Kent, OH 44242.

DOI:10.1523/JNEUROSCI.0270-10.2010

Copyright © 2010 the authors 0270-6474/10/3013955-11\$15.00/0

PKC-stimulated trafficking of GABA $_A$ receptors could contribute to this phenotype, we hypothesized that PKC ϵ regulates GABA $_A$ receptor trafficking. Here, we demonstrate that PKC ϵ decreases the number of GABA $_A$ receptors at the cell surface by binding, phosphorylating, and activating NSF.

Materials and Methods

Animal care. *Prkce*^{-/-} mice were generated by homologous recombination in J1 embryonic stem cells (Khasar et al., 1999). Male and female *Prkce*^{+/-} mice were maintained on inbred 129S4 and C57BL/6J backgrounds and crossed to produce *Prkce*^{+/-} C57BL/6J \times 129S4 F₁ hybrid mice for breeding. These mice were intercrossed to generate F₂ hybrid *Prkce*^{+/+} and *Prkce*^{-/-} littermates for experiments. Mice were genotyped by PCR of tail DNA. All animals used were males between 2 and 4 months of age at the time of experimentation. All procedures were conducted in accordance with institutional Institutional Animal Care and Use Committee policies.

Reagents. Peptides were synthesized by AnaSpec. An affinity-purified, polyclonal rabbit anti-phospho-NSF (S460) antibody (1:50) was generated against the phosphopeptide 455RHIKA[pS]TKVEV465 by ProSci. The ATP analog-specific kinase inhibitor 1-naphthyl-4-amino-5-(4-methylphenyl)-7-(*t*-butyl)pyrazolo[3,4-*d*]pyrimidine (1Na-PP1) was synthesized as described previously (Bishop et al., 1999). The MyrPalm-CFP expression plasmid was kindly provided by A. Newton (University of California, San Diego, La Jolla, CA). 6,7-Dinitroquinoxaline-2,3(1*H*,4*H*)-dione (DNQX) and *D*(-)-2-amino-5-phosphonopentanoic acid (*D*-APV) were purchased from Tocris Bioscience. Other reagents were from Sigma-Aldrich or from the sources listed below.

Cell culture. HEK293 cells stably expressing rat α 1 β 2 γ 2S GABA $_A$ receptors and the human analog-specific PKC ϵ mutant (AS-PKC ϵ , PKC ϵ M486A) were generated as described previously (Qi et al., 2007). Cells were maintained in minimal essential medium with 10% fetal bovine serum, 0.2 mM L-glutamine, 200 μ g/ml Geneticin (G418), 100 μ g/ml streptomycin, and 100 U/ml penicillin at 37°C in a humidified atmosphere of 5% CO₂ and 95% air. Primary cultures of hippocampal neurons were prepared from postnatal day 1 (P1) to P3 mice and cultured at 10⁵ cells per 200 mm² in Neurobasal A medium containing B27 and Glutamax (Invitrogen) at 37°C in a humidified atmosphere of 6% CO₂ and 94% air. One-half of the medium was changed the next day and every 7 d thereafter for up to 3 weeks.

Cell surface biotinylation. HEKGR-AS-PKC ϵ cells were grown on 100 mm poly-L-ornithine (100 μ g/ml in 15 mM borate buffer)-coated tissue culture dishes to 80–90% confluence. Cells were rinsed twice at 37°C with buffer containing 145 mM NaCl, 5 mM KCl, 10 mM glucose, 1 mM MgCl₂, and 1 mM CaCl₂ (adjusted with Tris base to pH 7.5), and incubated at 37°C with drugs in 10 ml of the same buffer. Cells were then rinsed with ice-cold PBS and cell surface proteins biotinylated using 1.5 mg/ml sulfo-NHS biotin (Pierce) as described previously (Qian et al., 1997). Biotinylated proteins were recovered from protein lysates on Immunopure Immobilized Monomeric Avidin beads (Pierce) and separated by SDS-PAGE on duplicate NuPAGE 4–12% Bis-Tris gels (Invitrogen). Proteins in one gel were transferred to nitrocellulose and subjected to Western blot analysis with rabbit anti-GABA $_A$ γ 2 antibody (1:1000; Alpha Diagnostics International) and HRP-conjugated goat anti-rabbit IgG (1:1000; Millipore Bioscience Research Reagents). Immunoreactive bands were visualized using enhanced chemiluminescence (Pierce) and quantified by scanning densitometry with ImageJ (<http://rsbweb.nih.gov/ij/>). To control for recovery of biotinylated proteins, we stained the duplicate gel with Coomassie blue and quantified 45–55 kDa proteins by scanning densitometry, and then used these results to normalize Western blot results for each corresponding biotinylated sample. Cell lysates were examined by Western blot analysis to detect levels of total γ 2 subunits. Values for γ 2 immunoreactivity were normalized for GAPDH immunoreactivity detected in the same samples using rabbit anti-GAPDH antibody (1:2000; Abcam).

Cell electrophysiology. HEK293 cells were plated on 35 mm plastic dishes coated with poly-D-lysine. Conventional whole-cell recordings were made with recording electrodes (5–8 M Ω) filled with the following (in mM): 145 *N*-methyl-D-glucamine (NMDG)-Cl, 2 MgCl₂, 0.1 CaCl₂, 5

EGTA, 10 HEPES, and 2 Mg²⁺-ATP, pH 7.3, adjusted with HCl. Cells were continuously perfused with an external solution (in mM: 145 NaCl, 3 KCl, 1.5 CaCl₂, 1 MgCl₂, 10 HEPES, and 10 *D*-glucose, pH 7.4, adjusted with NaOH). Cells were voltage clamped at -75 mV and whole-cell current was elicited by increasing concentrations of GABA that were fast applied for 5 s. There was a 1 min washout period between each GABA application with 1Na-PP1 or vehicle present in the washout solution. All recordings were obtained at room temperature. Data were expressed as current density by dividing whole-cell current by whole-cell capacitance and averaged to generate sigmoid dose–response curves for each treatment condition. For mouse hippocampal neurons (6–12 d *in vitro*), recording electrodes (5–8 M Ω) contained the following (in mM): 145 NMDG-Cl, 1 MgCl₂, 5 HEPES, 4 Mg²⁺-ATP, 2 Na⁺-ATP, adjusted to pH 7.3 with HCl. The external control solution used to continuously perfuse the neurons contained the following (in mM): 142 NaCl, 3 KCl, 1.5 CaCl₂, 1 MgCl₂, 10 HEPES, 10 *D*-glucose, adjusted to pH 7.4 with NaOH. The GABA dose–response relationship was found to be similar for *Prkce*^{+/+} and *Prkce*^{-/-} neurons, and a sub-EC₅₀ concentration of 10 μ M was chosen for current stability experiments during which GABA was applied for 5 s every 2.5 min for up to 40 min. All recordings were made at 33°C with cells voltage clamped at -75 mV. The ψ E RACK peptide (HDAPIGYD; 5 μ M) (Chen et al., 2001) was added to the pipette solution to activate PKC ϵ . GABA and 1Na-PP1 were applied by local perfusion, and whole-cell currents were filtered at 2 kHz and digitized at 5 kHz. Where used, the final concentration of DMSO applied to neurons was <0.1% (v/v), which had no effect on GABA $_A$ receptor currents.

Slice electrophysiology. Male mice (P45–P65) were anesthetized with halothane and then decapitated. Their brains were rapidly removed and put in ice-cold artificial CSF (ACSF) containing the following (in mM): 124 NaCl, 26 NaHCO₃, 2 KCl, 2 CaCl₂, 2 MgCl₂, 1.25 KH₂PO₄, 10 glucose, equilibrated with 95% O₂ and 5% CO₂. Horizontal hippocampal sections (300 μ m) were cut with a vibratome in ice-cold ACSF. Slices were left to recover at 30°C for 30 min before transfer to a recording chamber at room temperature. Slices were continuously superfused with ACSF at 2 ml/min at room temperature. Whole-cell recordings of CA1 hippocampal pyramidal neurons were obtained with electrodes that had a resistance of 5–8 M Ω when filled with pipette solution containing the following (in mM): 145 K-gluconate, 2 MgCl₂, 10 HEPES, 0.5 EGTA, 2 Mg-ATP, Na-GTP, and 5 phosphocreatine, pH 7.3, adjusted with KOH; osmolarity, 270–280 mOsm. Cells were held under voltage clamp at -45 mV, and responses were low-pass filtered at 2 kHz, and digitized at 10 kHz. Synaptic responses were evoked by monophasic current pulses (200 μ s; 30 μ A; 4 Hz) delivered by a pair of monopolar tungsten electrodes placed in CA1 stratum radiatum. Responses mediated by GABA $_A$ receptors were pharmacologically isolated by adding 10 μ M DNQX and 30 μ M *D*-APV to the ACSF to block glutamate receptors. Addition of bicuculline abolished these currents indicating that they were generated by GABA $_A$ receptors. Recordings began immediately after establishing the whole-cell configuration. Only recordings that showed a <30% change in series resistance over the 20–40 min recording period were analyzed.

Immunoprecipitation. HEKGR-AS-PKC ϵ cells were cultured as described above for biotinylation studies. Cells were collected in buffer containing 50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 5 mM EDTA, 5 mM EGTA, 1% NP-40, and Complete protease inhibitor mixture (Roche Applied Science) and lysed by freezing and thawing. Lysates were centrifuged at 20,817 \times g for 15 min at 4°C. Whole forebrains from 2- to 4-month-old mice were homogenized using 15 strokes of a Teflon-glass homogenizer in ice-cold homogenization buffer containing 25 mM HEPES, pH 7.4, 150 mM NaCl, 1% Triton X-100, 1 mM EDTA, 0.5 mM ATP, Complete protease inhibitor mixture (Roche Applied Science), phosphatase inhibitor mixture 1 (Sigma-Aldrich), and 1 mM PMSF. The homogenate was centrifuged at 10,000 \times g for 10 min at 4°C. Supernatants (500 μ g protein in 1 ml buffer) were incubated overnight at 4°C with rabbit anti-NSF antibody (5 μ g/ml; Millipore Biotechnology) and protein A-agarose (50 μ l; Roche Applied Science) and separated on NuPAGE 4–12% Bis-Tris gels (Invitrogen) in MOPS (4-morpholinepropane-sulfonic acid) running buffer. Proteins were examined by Western blot analysis with goat anti-PKC ϵ (1:1000; Santa Cruz Biotechnology), mouse anti-NSF antibody (1:1000; BD Biosciences), or anti-GABA $_A$ receptor γ 2 subunit (1:500; PhosphoSolutions). Phosphorylation of NSF at S460 was




Figure 1. PKC ϵ regulates cell surface levels of GABA $_A$ receptors. **A**, HEKGR-AS-PKC ϵ cells were incubated for 1 h with 0.03% DMSO vehicle (con) ($n = 10$), 1 μ M 1Na-PP1 ($n = 10$), 30 nM PMA ($n = 5$), or 1Na-PP1 plus PMA ($n = 4$). The top panel is a representative Western blot of biotinylated proteins detected using anti- γ 2-subunit antibodies. * $p < 0.05$ compared with untreated control (100%) by one-sample t tests. **B**, GABA-stimulated currents recorded from HEKGR-AS-PKC ϵ cells pretreated for 1 h with 0.03% DMSO vehicle (control; $n = 16$) or 3 μ M 1Na-PP1 ($n = 14$). 1Na-PP1 increased the GABA current density from 38.8 ± 3.8 to 54.7 ± 4.4 pA/pF ($p = 0.0086$) without altering the EC $_{50}$ for GABA, which was 13.8μ M (log EC $_{50} = -4.86 \pm 0.25$) in control and 8.7μ M (log EC $_{50} = -5.06 \pm 0.23$) in 1Na-PP1-treated cells ($p = 0.554$). **C**, GABA-gated currents recorded every 2.5 min for 30 min from cultured *Prkce*^{+/+} hippocampal neurons treated with buffer (control; $n = 6$) or with the PKC ϵ activator 5 μ M $\psi\epsilon$ RACK ($n = 10$), and from *Prkce*^{-/-} neurons treated with $\psi\epsilon$ RACK ($n = 6$). **D**, Normalized amplitudes at each time point were expressed relative to current measured at time 0. Compared with vehicle-treated *Prkce*^{+/+} cells (control; $n = 4$) treatment of *Prkce*^{+/+} neurons with $\psi\epsilon$ RACK ($n = 5$) decreased current amplitude, whereas in *Prkce*^{-/-} neurons ($n = 4$) $\psi\epsilon$ RACK had no effect ($F_{\text{time}(12,120)} = 11.59, p < 0.0001$; $F_{\text{condition}(2,120)} = 20.05, p = 0.0003$; $F_{\text{time by condition}(24,120)} = 6.47, p < 0.0001$). * $p < 0.05$ versus control by Bonferroni's *post hoc* test. **E**, Representative eIPSC traces recorded from hippocampal pyramidal neurons in acute brain slices treated with buffer (control), 5 μ M $\psi\epsilon$ RACK, or 5 μ M scrambled $\psi\epsilon$ RACK peptide. **F**, Normalized amplitudes of eIPSCs averaged every 3 min and expressed relative to current measured immediately after establishing whole-cell patch configuration at time = 0. Compared with control cells ($n = 5$), treatment with $\psi\epsilon$ RACK ($n = 5$) decreased eIPSC amplitude over the 30 min test period, whereas scrambled $\psi\epsilon$ RACK ($n = 4$) had no effect ($F_{\text{time}(12,132)} = 6.85, p < 0.0001$; $F_{\text{treatment}(2,132)} = 23.41, p = 0.0001$; $F_{\text{time by treatment}(24,132)} = 5.73, p < 0.0001$). * $p < 0.05$ versus control by Bonferroni's *post hoc* test. Error bars indicate SEM.

detected using polyclonal rabbit anti-NSF (S460) antibody (diluted 1:50). Immunoreactivity was detected using appropriate HRP-conjugated anti-IgG secondary antibodies (1:1000; Jackson ImmunoResearch) followed by enhanced chemiluminescence (Pierce).

Pull-down assay. FLAG-tagged human PKC ϵ (in pcDNA3 from A. Toker, Harvard University, Cambridge, MA) was purified from COS-7 cells using the FLAG system (Sigma-Aldrich) (Qi et al., 2007). FLAG-

human PKC ϵ was immobilized on anti-FLAG antibody M2 conjugated to agarose (50 μ l; Sigma-Aldrich), and then incubated for 2 h at 4°C with 0, 2.5, or 5 μ g of recombinant His $_6$ -NSF in binding buffer (10 mM Tris-HCl, pH 7.5, 300 mM NaCl, 2 mM EDTA, 0.5 mM ATP). After centrifugation at $20,817 \times g$ for 5 min, beads were washed three times with binding buffer. Bound proteins were separated in NuPAGE Bis-Tris gels (Invitrogen) and detected by Western blot analysis with monoclonal anti-NSF antibody (1:1000; BD Biosciences) and polyclonal anti-PKC ϵ antibody SN134 (1:1000) (Qi et al., 2007). Immunoreactivity was quantified by scanning densitometry using ImageJ, and results for NSF immunoreactivity were normalized to PKC ϵ immunoreactivity.

Gel overlay assay. For overlay assays, NSF (0.5 μ g) was run on NuPAGE Bis-Tris gels (Invitrogen) and transferred to nitrocellulose Hybond C Extra membranes (GE Healthcare). Membranes were incubated in blocking buffer containing TBS-Tween 20 and 2% nonfat milk for 1 h at room temperature and incubated overnight at 4°C with 10 nM recombinant PKC ϵ (Invitrogen) in overlay buffer containing 50 mM Tris-HCl, pH 7.5, 10 mM β -mercaptoethanol, 200 mM NaCl, 1 mM EGTA, 0.1% BSA, 1% polyethylene glycol, 0.3 μ g/ μ l L- α -phosphatidylserine, 62 μ g/ml 1,2-diolein, and 0.03% Triton X-100. After being washed for 10 min three times in TBS-Tween 20 containing 1% nonfat milk, the membranes were incubated at room temperature for 4 h with anti-PKC ϵ antibody SN134 (1:1000) in TBS-Tween 20 containing 1% nonfat milk. Membranes were then washed for 10 min with TBS-Tween 20 four times, and PKC ϵ bound to membranes was detected by HRP-conjugated anti-rabbit IgG and enhanced chemiluminescence. The amount of mutant NSF loaded in each lane was detected by anti-NSF antibody (1:5000; BD Biosciences), HRP-conjugated donkey anti-mouse IgG antibodies (1:1000; Jackson ImmunoResearch), and enhanced chemiluminescence.

Immunofluorescence microscopy. HEKGR-AS-PKC ϵ cells were plated on poly-D-lysine (BD Biosciences)-coated chamber slides at 5×10^3 cells per 200 mm 2 and grown overnight in minimal essential medium with 10% fetal bovine serum, 0.2 mM L- α -glutamine, 200 μ g/ml G418, 100 μ g/ml streptomycin, and 100 U/ml penicillin at 37°C in a humidified atmosphere of 5% CO $_2$ and 95% air. To identify plasma membrane, we transfected these cells with MyrPalm-CFP, which encodes cyan fluorescent protein fused to the myristoylation and palmitoylation sequence of Lyn kinase (Violin et al., 2003), using the SuperFect Transfection Reagent (QIAGEN). Cells were used for experiments 24 h later. After drug treatment, cells were fixed for 15 min with 4% paraformaldehyde in PBS, permeabilized for 5 min with 0.1% Triton X-100 in PBS, and blocked for 1 h in 10% normal donkey serum (NDS) and 0.2% BSA in PBS. Samples were incubated overnight at 4°C with the following primary antibodies diluted in PBS containing 2% NDS and 0.2% BSA: chicken anti-GFP (green fluorescent protein) (1:100; Invitrogen), mouse anti-NSF (1:200; BD Biosciences), and goat anti-PKC ϵ (1:100; Santa Cruz

Biotechnology). After three washes with PBS, cells were incubated with donkey DyLight 649-conjugated anti-goat, DyLight 549-conjugated anti-mouse, and DyLight 488-conjugated anti-chicken IgY secondary antibodies (1:200; Jackson ImmunoResearch).

Mouse hippocampal neurons were plated on poly-D-lysine (BD Biosciences)-coated chamber slides at 10^5 cells per 200 mm^2 for 14–21 d before treatment with drugs. To detect GABA $_A$ receptors, live cultures were incubated with rabbit anti-GABA $_A$ receptor $\gamma 2$ subunit antibody (1:100; Alomone Labs) diluted in Ringer's solution containing $0.5\ \mu\text{M}$ tetrodotoxin for 90 min at room temperature. After three 10 min washes in Ringer's solution, neurons were fixed with methanol for 10 min at -20°C , washed with PBS, and incubated for 90 min at room temperature with mouse anti-gephyrin (1:200; Synaptic Systems) in PBS containing 10% NDS. To detect NSF and gephyrin, methanol-fixed cultures were incubated with rabbit anti-NSF (1:200; Millipore Biotechnology) and mouse anti-gephyrin antibodies. After three washes with PBS, cells were incubated with DyLight 549-conjugated anti-rabbit and DyLight 488-conjugated anti-mouse IgG secondary antibodies (1:200; Jackson ImmunoResearch).

Immunofluorescence was detected using a Zeiss LSM 510 confocal microscope (Carl Zeiss MicroImaging) with a Plan ApoChromat 63 \times /1.40 numerical aperture oil-immersion objective. Quantification of colocalization was performed using ImageJ (Abramoff et al., 2004) with the PSC colocalization plug-in (French et al., 2008). For studies using HEK293 cells, at least 26 cells were analyzed for each treatment condition. For experiments with neurons, neurites were analyzed in at least seven images (1024×1024 pixels; 7 pixels/ μm) for each treatment condition. Results are presented as Spearman's rank correlation coefficients. Unlike the Pearson's product-moment correlation coefficient, which measures the strength of the linear relationship between two signals, the Spearman's coefficient can detect linear and nonlinear relationships (French et al., 2008). The analysis gives values from -1 (a complete negative correlation) to $+1$ (a complete positive correlation), with zero representing no correlation. Nontransfected HEK293 cells and gephyrin-negative neurites were excluded from the analyses. Results were analyzed using Mann-Whitney tests and differences between median values were considered significant when $p < 0.05$.

In vitro kinase assay. In vitro phosphorylation of NSF or GST- $\gamma 2\text{S}$ by PKC ϵ was performed using $0.07\ \mu\text{M}$ FLAG-tagged human PKC ϵ purified from COS-7 cells (Qi et al., 2007) in $10\ \mu\text{l}$ of kinase buffer containing 20 mM HEPES, pH 7.4, 0.1 mM EGTA, 0.03% Triton X-100, 10 mM MgCl $_2$, 0.48 $\mu\text{g}/\mu\text{l}$ L- α -phosphatidylserine (Avanti Polar Lipids), 1 μM phorbol 12-myristate, 13-acetate (PMA), and 0.5 mM ATP. The kinase reaction was initiated at 37°C by adding 10.8 pmol of recombinant NSF or 18.8 pmol of GST- $\gamma 2\text{S}$ and $10\ \mu\text{Ci}$ of [γ - ^{32}P]ATP. At different time points, $10\ \mu\text{l}$ aliquots of the reaction mixture were removed and added to $2.5\ \mu\text{l}$ of $5\times$ SDS buffer to stop the reaction. Proteins were separated on NuPAGE Bis-Tris gels (Invitrogen) and stained using SimplyBlue (Invitrogen). Phosphorylated NSF was detected by phosphorimaging. The specific radioactivity per mole of [γ - ^{32}P]ATP was determined by counting diluted aliquots of the stock [γ - ^{32}P]ATP solution to obtain a conversion value for calculating the molar ratio of ^{32}P to NSF. The stoichiometry of phosphate incorporation into NSF was expressed as moles of incorporated γ - ^{32}P per moles of NSF.




Figure 2. PKC ϵ interacts directly with NSF in a complex containing GABA $_A$ receptor $\gamma 2$ subunits. **A**, The left panel is a representative experiment showing proteins immunoprecipitated from HEKGR-AS-PKC ϵ cell lysates with anti-NSF antibody and detected by Western blot analysis with anti-NSF, anti-PKC ϵ , or anti- $\gamma 2$ -subunit antibodies. Data are quantified in the right panel, showing that, after 45 min of treatment, $1\ \mu\text{M}$ 1Na-PP1 decreased and $1\ \mu\text{M}$ PMA increased the amount of PKC ϵ coimmunoprecipitated with NSF [$n = 3$; $*p < 0.05$ compared with control (con) and $^\dagger p < 0.05$ compared with PMA by Newman-Keuls *post hoc* test]. **B**, Representative Western blots with anti-NSF, anti-PKC ϵ , and anti- $\gamma 2$ -subunit antibodies of brain lysates from *Prkce* $^{+/+}$ and *Prkce* $^{-/-}$ mice after immunoprecipitation with anti-NSF antibody. **C**, The top panel shows representative Western blots from a pull-down assay using recombinant NSF and purified FLAG-PKC ϵ ($2\ \mu\text{g}$) immobilized on anti-FLAG antibody-conjugated agarose. The amount of NSF pulled down with immobilized PKC ϵ increased linearly with the amount of NSF added to the assay (bottom panel; $n = 4$). **D**, Representative overlay assay showing Ponceau S-stained-NSF and GST immobilized on a nitrocellulose membrane after SDS-PAGE (Ponceau S; left) and PKC ϵ immunoreactivity detected using an anti-PKC ϵ antibody (PKC ϵ overlay; right) after incubation of the membrane with recombinant PKC ϵ . This experiment was repeated three times with similar results. Error bars indicate SEM.

Purification of NSF. Recombinant His-tagged Chinese hamster ovary NSF cDNA in the vector pQE9 was kindly provided by Dr. Sidney Whiteheart (University of Kentucky, Lexington, KY). Recombinant His-NSF and its mutants were isolated from *Escherichia coli* M15 by chromatography on nickel-agarose columns (QIAGEN). To preserve the enzymatic activity of NSF, 0.5 mM ATP were added to the lysis buffer and 0.5 mM ADP were added to the washing and elution buffers. Bacterial cultures were grown at 37°C in LB broth containing $100\ \mu\text{g}/\text{ml}$ ampicillin and $25\ \mu\text{g}/\text{ml}$ kanamycin until the OD $_{600}$ reached 0.6, and then induced with 1 mM IPTG (isopropyl β -D-thiogalactopyranoside) for 4 h. The cells were collected and resuspended in 25 ml of lysis buffer containing 20 mM HEPES, pH 8.0, 50 mM NaCl, 0.5 mM ATP, 5 mM MgCl $_2$, 5 mM β -mercaptoethanol, and 1 mM PMSF. Extracts were solubilized by sonication. The unsolubilized material was removed by centrifugation at $200,000 \times g$ for 30 min at 4°C . After affinity binding of the supernatant to nickel-agarose columns, the columns were washed once with lysis buffer containing 10 mM imidazole and once with lysis buffer containing 0.5 mM ADP and 20 mM imidazole. The proteins were eluted using 100 mM HEPES at pH 8.0, 300 mM NaCl, 10% glycerol, 0.5 mM ADP, and 300 mM imidazole buffer. After concentration in Centricon K50 columns, the proteins were dialyzed against buffer containing 20 mM HEPES at pH 7.4, 100 mM NaCl, 10% glycerol, 1 mM MgCl $_2$, 0.5 mM ADP, and 1 mM β -mercaptoethanol and stored at -80°C . The concentration of recombinant NSF was determined by comparison with bovine serum albumin standards on NuPAGE Bis-Tris gels (Invitrogen). The purity of the final proteins was $>95\%$, as assessed by Coomassie blue staining.

Mass spectrometry. NSF proteins were phosphorylated *in vitro* by PKC ϵ for 3 h as described above. The reaction mixtures were fractionated by SDS-




Figure 3. Membrane translocation of NSF in response to activation of PKC ϵ . **A**, HEK293T cells were transfected with MyrPalm-CFP and incubated for 60 min with control vehicle (0.1% DMSO; Con), 1 μ M 1Na-PP1, or 1 μ M PMA, or with 1Na-PP1 for 15 min before addition of PMA for another 60 min. CFP (green), NSF (red), and PKC ϵ (blue) immunoreactivity were detected by confocal microscopy. Merged images indicate colocalization of CFP and NSF (yellow) or CFP and PKC ϵ (cyan). PMA (1 μ M) induced translocation of NSF and PKC ϵ to cell membrane. **B**, **C**, The colocalization between these proteins was quantified and expressed as Spearman's rank correlation coefficients. The results indicate that 1 μ M PMA increased the colocalization of both NSF ($n = 31$ –57 cells) and PKC ϵ ($n = 26$ –38 cells) with CFP. * $p < 0.05$ compared with other conditions (Dunn's multiple-comparisons test). Scale bar, 10 μ m. Error bars indicate SEM.

PAGE and stained with Colloidal Blue (Invitrogen). Gel slices containing NSF were digested with trypsin or AspN protease. Samples from the digests were analyzed by nano-liquid chromatography–mass spectrometry/mass spectrometry (nano-LC-MS/MS) using a LC-Packings HPLC (Dionex) coupled to a QStar XL mass spectrometer (Applied Biosystems). Peptides were first desalted on a 300 μ m \times 5 mm PepMap C18 trap column with 0.1% formic acid in HPLC-grade water at a flow rate of 20 μ l/min. After being desalted for 6 min, peptides were backflushed onto a LC Packings 75 μ m \times 15 cm C18 nano column (3 μ m; 100 Å) at a flow rate of 200 nl/min. Peptides were eluted with a 30 min gradient of 3–40% acetonitrile in 0.1% formic acid. Mass ranges for the MS survey scan and MS/MS were m/z 300–1800 and m/z 50–1800, respectively. The scan times for MS and MS/MS were 1.0 and 2.0 s, respectively. The top three, multiply charged ions with MS peak intensity >30 counts/scan were chosen for MS/MS fragmentation with a precursor ion dynamic exclusion of 60 s. Data were searched against a home-built database that included the NSF sequence. Four modifications were included in the database search as follows: carbamidomethyl (C), oxidation (M), phospho (ST), and phospho (Y).

NSF peptide phosphorylation. The following peptides (based on amino acids 452–468 of NSF) were dissolved in ddH $_2$ O to prepare 1 mM stocks: native NSF (AMNRRHIKASTKVEVDME), S460A mutant (AMNRRHI-

KAATKVEVDME), T461A mutant (AMNRRHIKASAKVEVDME), and S460A/T461A double mutant (AMNRRHIKAAAKVEVDME). Human recombinant PKC ϵ (Invitrogen) was diluted in 0.1 mg/ml BSA, 0.05% Triton X-100 to 4 ng/ μ l. Phosphorylation was performed with 9.5 nM PKC ϵ and 100 μ M NSF peptide in 25 μ l of kinase buffer containing 20 mM Tris-HCl, pH 7.5, 10 mM MgCl $_2$, 1 μ M PMA, 0.48 mg/ml L- α -phosphatidylserine, 0.25 mM EGTA, 0.1 mg/ml BSA, 2.5 μ M ATP, and 0.5 μ Ci of [32 P]ATP. After 90 min incubation at 27°C, the reaction was terminated by adding 12.5 μ l of 7.5 M guanidine hydrochloride. Eight microliters of reaction mixture were spotted onto a streptavidin-coated SAM Biotin Capture membrane (Promega). The membrane was washed once with 2 M NaCl for 30 s, three times with 2 M NaCl for 2 min, four times with 2 M NaCl in 1% H $_3$ PO $_4$ for 2 min, and twice with double-distilled H $_2$ O for 30 s. The membrane was air dried at 27°C for 60 min, and radioactivity remaining on the membrane was measured by liquid scintillation counting.

ATPase assay. NSF (600 ng) was phosphorylated *in vitro* for 30 min using 200 ng of PKC ϵ as described above but in the presence of 1 mM ATP without [32 P]ATP. ATPase assays were then performed in 30 μ l of 25 mM Tris-HCl, pH 9.0, 100 mM KCl, 0.65 mM β -mercaptoethanol, 2 mM MgCl $_2$, 0.5 mM ATP, and 10% glycerol for 1 h at 37°C. The reaction was initiated by the addition of 10 μ Ci of [32 P]ATP. An aliquot (1 μ l) of the reaction mixture was separated on cellulose PEI thin-layer plates (J. T. Baker) in 0.8 M acetic acid and 0.8 M LiCl and analyzed using a Typhoon 9410 scanner (GE Healthcare). The ATPase activity of NSF was calculated as the percentage of released inorganic 32 P using the following formula: ATPase activity = 32 P/([32 P] + [32 P]ATP).

Cell surface ELISA. Hippocampal neurons were grown on four-well poly-D-lysine-coated slides. After drug treatment at 37°C in medium, slides were placed on ice and washed twice in ice-cold PBS, fixed for 10 min in ice-cold PBS containing 3% paraformaldehyde, and washed three times with PBS. Neurons were then incubated with rabbit anti-GABA $_A$ γ 2 (extracellular) antibody (1:250; Alomone Labs), followed by anti-rabbit IgG (H+L) (1:5000; Jackson ImmunoResearch) at room temperature for 1 h, washed six times with ice-cold PBS, and incubated with 0.5 ml per well of 3,3',5,5'-tetramethylbenzidine (Sigma-Aldrich) for 1 h at room temperature. The supernatant (100 μ l) was transferred to a 96-well plate, and absorbance was measured at 655 nm. Results were expressed as a percentage of absorbance detected in neurons treated with vehicle alone.

Statistical analysis. Quantitative data were expressed as mean \pm SEM values and analyzed using Prism 5.0 (GraphPad Software). Data were tested for normality by the Kolmogorov–Smirnov test and then for differences between means by *t* tests or one-way ANOVA with Newman–Keuls *post hoc* tests. Data that were not normally distributed were analyzed by Mann–Whitney tests, or by Kruskal–Wallis and *post hoc* Dunn's multiple-comparison tests. Dose–response curves were analyzed by nonlinear curve fitting. Values of $p < 0.05$ were considered to be statistically significant.

Results

PKC ϵ regulates the cell surface level of GABA $_A$ receptors

To examine the role of PKC ϵ in GABA $_A$ receptor trafficking, we investigated phorbol ester-stimulated downregulation of GABA $_A$

receptors. We used HEKGR-AS-PKC ϵ cells, which stably express $\alpha 1\beta 2\gamma 2$ GABA $_A$ receptors and an ATP analogue-specific mutant of PKC ϵ (AS-PKC ϵ) that can be selectively inhibited by the AS-kinase inhibitor 1Na-PP1 at concentrations that do not affect native kinases (Qi et al., 2007). We treated these cells with the PKC activator PMA, with or without 1Na-PP1, and measured the level of GABA $_A$ $\gamma 2$ subunits at the cell surface by Western blot analysis of biotinylated cell surface proteins (Fig. 1A). PMA (30 nM) reduced the cell surface level of $\gamma 2$ subunits by 43%, whereas 1 μ M 1Na-PP1 increased it by 33% and antagonized the effect of PMA. These treatments did not alter the total level of $\gamma 2$ subunits in the cell lysate (supplemental Fig. 1, available at www.jneurosci.org as supplemental material).

These findings suggested that PKC ϵ regulates the level of GABA $_A$ receptors at the cell surface. To investigate this hypothesis further, we measured GABA-stimulated currents in HEKGR-AS-PKC ϵ cells. Treatment with 3 μ M 1Na-PP1 increased the GABA current density by 41% without altering the EC $_{50}$ for GABA (Fig. 1B). This increase in GABA efficacy was similar in magnitude to the increase in cell surface $\gamma 2$ subunits induced by 1Na-PP1 (Fig. 1A), indicating that PKC ϵ regulates GABA efficacy by altering the number of functional GABA $_A$ receptors at the cell surface.

We next investigated whether PKC ϵ also regulates GABA $_A$ receptors in neurons by studying mouse hippocampal neurons from P1–P3 wild-type and *Prkce*^{-/-} mice cultured for 6–12 d *in vitro*. Treatment with 5 μ M ψ E RACK, a specific PKC ϵ activator (Chen et al., 2001) administered via the recording pipette, reduced GABA-stimulated currents in wild-type neurons but had no effect on *Prkce*^{-/-} neurons (Fig. 1C,D). In wild-type brain slices from P45–P65 mice, 5 μ M ψ E RACK administered in the recording pipette reduced electrically evoked, GABA-mediated inhibitory postsynaptic currents in hippocampal pyramidal neurons, whereas 5 μ M of a scrambled peptide had no effect (Fig. 1E,F). These results demonstrate that PKC ϵ downregulates native GABA $_A$ receptors in hippocampal neurons, which leads to decreased GABA-mediated inhibitory synaptic transmission.

PKC ϵ associates with NSF and GABA $_A$ $\gamma 2$ subunits

We next investigated mechanisms by which PKC ϵ regulates the number of GABA $_A$ receptors at the cell surface. Since PKC phosphorylation of GABA $_A$ receptor subunits does not regulate receptor trafficking (Chapell et al., 1998; Connolly et al., 1999), we considered as possible substrates proteins that associate with GABA $_A$ receptors and are known to regulate trafficking of other receptors. We focused attention on NSF since it regulates the trafficking of AMPA receptors (Collingridge et al., 2004). We found that NSF coimmunoprecipitated with both PKC ϵ and GABA $_A$ $\gamma 2$ subunits in lysates of HEKGR-AS-PKC ϵ cells (Fig. 2A). Incubation with 1 μ M PMA for 45 min increased the amount of coimmunoprecipitated PKC ϵ by ~60%, whereas 1Na-PP1 decreased it by 60% and prevented the effect of PMA ($F_{(3,8)} = 16.4$; $p = 0.0009$) (Fig. 2A). In contrast, the amount of GABA $_A$ $\gamma 2$ that coimmunoprecipitated with NSF was not altered by either treatment. The abundance of NSF, PKC ϵ , and $\gamma 2$ in the lysates was also not altered by these treatments (supplemental Fig. 2A, available at www.jneurosci.org as supplemental material).

To test whether PKC ϵ , NSF, and GABA $_A$ receptors also form a protein complex in the brain, we prepared whole-brain lysates from *Prkce*^{+/+} and *Prkce*^{-/-} mice. The abundance of NSF and GABA $_A$ $\gamma 2$ subunits was similar in *Prkce*^{+/+} and *Prkce*^{-/-} lysates (supplemental Fig. 2B, available at www.jneurosci.org as supplemental material). NSF could be coimmunoprecipitated with




Figure 4. PKC activation alters the colocalization of NSF and GABA $_A$ receptors with gephyrin. **A**, PMA increases the colocalization of NSF with gephyrin in cultured hippocampal neurons. Neurons were incubated with the control vehicle (0.02% DMSO; $n = 14$ images) or 200 nM PMA ($n = 14$). NSF (green) and gephyrin (red) were detected by confocal microscopy. Representative neurites in a $250 \times 500 \mu\text{m}$ region are shown. Merged images show colocalization of NSF with gephyrin in yellow. **B**, Compared with control (0.02% DMSO; $n = 7$ images), 200 nM PMA ($n = 15$) decreased the colocalization of GABA $_A$ $\gamma 2$ subunits with gephyrin. Merged images show colocalization of $\gamma 2$ subunits (green) and gephyrin (red) in yellow. Higher magnification merged images are shown in the bottom panels. Scale bars, 10 μm .

PKC ϵ and GABA $_A$ $\gamma 2$ subunits in lysates from *Prkce*^{+/+} mice (Fig. 2B). The amount of GABA $_A$ $\gamma 2$ coimmunoprecipitated with NSF from *Prkce*^{-/-} brain lysates was $97 \pm 3\%$ of the amount coimmunoprecipitated from *Prkce*^{+/+} lysates ($n = 3$ for each genotype; $p = 0.45$), indicating that the association between NSF and GABA $_A$ $\gamma 2$ subunits occurs independently of PKC ϵ . Reciprocal immunoprecipitations using antibodies against PKC ϵ or GABA $_A$ $\gamma 2$ confirmed their association with NSF in HEKGR-AS-PKC ϵ cell and *Prkce*^{+/+} brain lysates (supplemental Fig. 3, available at www.jneurosci.org as supplemental material).

NSF and PKC ϵ may interact directly or require another protein to bind in a complex. To investigate these possibilities, we first performed a pull-down assay using FLAG epitope-tagged PKC ϵ immobilized on agarose beads and His-tagged NSF (Fig. 2C). There was a linear correlation between the amount of NSF immunoreactivity recovered from the PKC ϵ -agarose beads and the amount of His-tagged NSF that was added to the assay ($r_p = 0.995$; $p = 0.0204$; $n = 4$). We also performed a gel overlay assay using NSF immobilized on nitrocellulose membranes (Fig. 2D). Recombinant PKC ϵ bound selectively to immobilized NSF but not to immobilized GST (Fig. 2D). These findings indicate that NSF and PKC ϵ can bind directly to each other.




Figure 5. PKCε phosphorylates NSF at S460 and regulates NSF ATPase activity. **A**, Representative autoradiograph of phosphorylated NSF (top panel) and a scanned image of a Coomassie blue-stained gel (bottom panel). **B**, *In vitro* phosphorylation of NSF peptides by PKCε is reduced by alanine mutations at S460 and T461 ($n = 3$ each). $*p < 0.05$ compared with native NSF by Newman–Keuls *post hoc* tests. **C**, Representative overlay assay showing NSF mutants immobilized on a nitrocellulose membrane after SDS-PAGE with PKCε immunoreactivity detected using an anti-PKCε antibody (PKCε overlay) after incubation of the membrane with recombinant PKCε. PKCε binding to alanine mutants (S460A, T461A) was reduced while binding to glutamate mutants (S460E, T461E) was increased ($p < 0.05$, one-sample *t* tests; $n = 5$). **D**, Representative autoradiograph of a thin-layer chromatogram showing increased NSF ATPase activity after PKCε phosphorylation of NSF. Radiolabeled ATP and released inorganic phosphate (Pi) are indicated by arrows. **E**, PKCε phosphorylation increased NSF ATPase activity and NEM blocked both baseline and PKCε-stimulated activity ($n = 3$). $*p < 0.05$ compared with NSF and $†p < 0.05$ compared with NSF + PKCε but not NSF + NEM by Newman–Keuls *post hoc* tests. **F**, The S460A NSF mutation prevented enhancement of ATPase activity by PKCε ($n = 3$). $*p < 0.05$ compared with the corresponding untreated control (100%) by one-sample *t* tests. Error bars indicate SEM.

Activation of PKCε causes NSF to translocate to the plasma membrane

Since phorbol esters cause translocation of PKC to the plasma membrane (Song and Messing, 2005) and our results indicated that NSF interacts with activated PKCε, we predicted that PMA would also cause NSF to translocate to the plasma membrane. We

examined this possibility by performing immunofluorescence studies in intact HEKGR-AS-PKCε cells using MyrPalm-CFP as a membrane marker (Violin et al., 2003). Treatment with PMA (1 μM) for 60 min increased the colocalization of NSF with MyrPalm-CFP, but addition of 1 μM 1Na-PP1 for 15 min before PMA blocked colocalization of NSF and MyrPalm-CFP (Fig. 3A, B) ($p < 0.0001$, Kruskal–Wallis test). PMA also increased the colocalization of PKCε and MyrPalm-CFP, and this effect was reduced by pretreatment with 1Na-PP1 (Fig. 3A, C) ($p < 0.0001$, Kruskal–Wallis test). These results indicate that phorbol esters induce a translocation of NSF to the plasma membrane that depends on PKCε activity.

We next investigated the association of NSF and GABA_A receptors with gephyrin, which is involved in the synaptic localization of GABA_A receptors (Essrich et al., 1998). We examined neurites of primary hippocampal neurons cultured from P1–P3 mice for 14–21 d *in vitro* and found that 200 nM PMA increased the colocalization of NSF and gephyrin ($p < 0.0001$), increasing the Spearman’s coefficient from 0.467 ± 0.008 in cells treated with the 0.02% DMSO vehicle ($n = 14$) to 0.538 ± 0.010 in cells treated with PMA ($n = 14$) (Fig. 4A). In contrast, PMA reduced colocalization of gephyrin and GABA_A γ2 subunits ($p = 0.0012$), decreasing the Spearman’s coefficient from 0.547 ± 0.017 in DMSO-treated cultures ($n = 7$) to 0.374 ± 0.030 in PMA-treated cultures ($n = 15$) (Fig. 4B). These results suggest that PKC activation recruits NSF to inhibitory synapses while downregulating the number of GABA_A receptors.

PKCε phosphorylation enhances the ATPase activity of NSF

Since PKCε and NSF interact, we next tested whether NSF is a PKCε substrate. *In vitro* kinase assays showed that PKCε phosphorylates NSF at a rate similar to the major intracellular loop of the GABA_A γ2S subunit (GST-γ2S), which contains a PKCε phosphorylation site at S327 (Fig. 5A) (Qi et al., 2007). PKCε phosphorylated NSF to a maximal stoichiometry of 0.917 ± 0.069 ($n = 3$). Nano-LC-MS/MS of *in vitro* phosphorylated and trypsin-digested NSF identified a single phosphorylated peptide that contains one serine and one threonine residue (underlined) (ASTKVEVD-MEK; amino acids 459–469) (supplemental Fig. 4, available at www.jneurosci.org as supplemental material). To test whether PKCε phosphorylates either site, we generated NSF peptides containing S460A, T461A, or both mutations. PKCε phosphorylated an NSF peptide containing the native sequence, but showed re-




Figure 6. PKC ϵ phosphorylates NSF *in vivo*. **A**, An affinity-purified rabbit anti-phospho-NSF (S460) antibody was generated that detects recombinant NSF phosphorylated by PKC ϵ (P), but not unphosphorylated NSF (N). The antibody was not immunoreactive against the NSF S460A mutant before or after incubation with PKC ϵ in a kinase reaction. **B**, Representative Western blot showing that the anti-phospho-NSF (S460) antibody detected less immunoreactivity in brain lysates from *Prkce*^{-/-} mice than in lysates from wild-type littermates. **C**, The ratio of P-NSF (S460) to NSF immunoreactivity was reduced in samples from *Prkce*^{-/-} ($n = 4$) compared with *Prkce*^{+/+} ($n = 4$) mice (* $p < 0.05$, two-tailed t test). Error bars indicate SEM.

duced ability to phosphorylate peptides containing either mutation ($F_{(3,8)} = 85.54$; $p < 0.0001$) (Fig. 5B). These results indicate that PKC ϵ phosphorylates NSF at both S460 and T461 *in vitro*.

To determine whether PKC ϵ phosphorylation of NSF regulates NSF function, we first examined the effect of phosphorylation on the interaction of NSF with PKC ϵ using gel overlay assays. Mutation of S460 or T461 to alanine disrupted binding to PKC ϵ (Fig. 5C). Conversely, mimicking phosphorylation by glutamic acid substitution at these residues facilitated binding to PKC ϵ . The results indicate that phosphorylation of NSF at S460 and T461 is necessary for its interaction with PKC ϵ .

We next investigated whether PKC ϵ phosphorylation regulates the enzymatic activity of NSF. Phosphorylation of NSF by PKC ϵ increased NSF ATPase activity by 50% ($F_{(3,8)} = 17.34$; $p = 0.0007$) (Fig. 5D,E). To ensure that this increase in activity, measured as hydrolysis of ATP, was attributable to NSF and not to PKC ϵ , we treated samples with the NSF inhibitor *N*-ethylmaleimide (NEM). This treatment abolished nearly all ATPase activity, indicating that the ATPase activity measured in this assay was attributable to NSF. To determine whether S460 and T461 are both necessary for regulation of NSF by PKC ϵ , we examined NSF variants containing these mutations and found that only the S460A mutation prevented enhancement of NSF ATPase activity by PKC ϵ (Fig. 5F). These results indicate that PKC ϵ phosphorylation of NSF at S460 increases the catalytic activity of NSF.

Since these *in vitro* studies showed that S460 is involved not only in enhancing NSF ATPase activity but also in the interaction of NSF with PKC ϵ , we investigated whether PKC ϵ phosphorylates S460 *in vivo*. We used an affinity-purified, phosphospecific antibody raised against a peptide containing phospho-S460. This anti-phospho-NSF (S460) antibody detected recombinant NSF phosphorylated by PKC ϵ *in vitro*, but not the nonphosphorylated protein (Fig. 6A). Anti-phospho-NSF (S460) immunoreactivity was decreased by 35% in *Prkce*^{-/-} brain lysates compared with wild-type tissue (Fig. 6B,C), consistent with the prediction that PKC ϵ directly phosphorylates brain NSF at S460 *in vivo*.

NSF regulates the level of GABA $_A$ receptors at the cell surface

To determine whether NSF ATPase activity regulates the cell surface level of GABA $_A$ receptors, we used a peptide inhibitor of NSF (NSFi) (Morrell et al., 2005), which consists of an NSF oligomerization domain attached to a protein transduction domain from the human immunodeficiency virus transactivator of transcription (TAT) protein. NSFi penetrates cells and blocks NSF ATPase activity (Morrell et al., 2005). We hypothesized that, if NSF acts downstream of PKC ϵ , then inhibiting NSF function should increase cell surface levels of GABA $_A$ receptors and prevent PKC ϵ -induced downregulation of GABA $_A$ receptors. We first examined HEKGR-AS-PKC ϵ cells and found that treatment for 1 h with 30 nM PMA reduced cell surface GABA $_A$ $\gamma 2$ immunoreactivity by 35% (Fig. 7A), as expected (Fig. 1A). Treatment for 75 min with 1 μ M NSFi increased cell surface GABA $_A$ $\gamma 2$ by 52% and preincubation with 1 μ M NSFi for 15 min before addition of 30 nM PMA prevented the effect of PMA (Fig. 7A). A control TAT peptide had no effect compared with the DMSO (0.02%) vehicle-treated control. PMA and NSFi did not alter the level of total $\gamma 2$ immunoreactivity in cell lysates (supplemental Fig. 5A, available at www.jneurosci.org as supplemental material).

We next determined whether NSFi similarly regulates GABA $_A$ receptors in neurons. For these studies, we used an ELISA that allowed us to detect $\gamma 2$ subunits using fewer cells than required by the biotinylation assay. Treatment of hippocampal neurons from *Prkce*^{+/+} mice with 1 μ M ψ εRACK for 1 h decreased cell surface $\gamma 2$ subunits by 26%, whereas treatment for 75 min with 1 μ M NSFi increased them by 60% and preincubation with 1 μ M NSFi for 15 min before addition of 1 μ M ψ εRACK completely prevented downregulation by ψ εRACK (Fig. 7B). Control scrambled versions of both peptides had no effect compared with untreated control cells. ψ εRACK and NSFi did not alter the level of total $\gamma 2$ immunoreactivity in cell lysates (supplemental Fig. 5B, available at www.jneurosci.org as supplemental material).

Based on these findings, we predicted that inhibiting NSF would increase GABA $_A$ currents. To test this prediction, we examined eIPSCs in hippocampal slices (Fig. 7C,D) while administering 2 μ M NSFi, 5 μ M ψ εRACK, or both peptides in the internal recording solution. We found that ψ εRACK decreased evoked GABA $_A$ currents in CA1 neurons by ~62%, whereas NSFi increased them by 98% and inhibited the effect of ψ εRACK ($F_{(2,17)} = 11.75$; $p = 0.0009$). To ensure that NSFi acted postsynaptically, we administered it together with ψ εRACK through the patch pipette. As a result, NSFi was less effective in reversing the effect of ψ εRACK in hippocampal slices than in neuronal cultures (Fig. 7B), since neurons in culture could be preincubated with NSFi for 15 min before addition of ψ εRACK to the medium. These results in hippocampal slices demonstrate that inhibiting NSF increases GABA $_A$ currents at inhibitory synapses on hippocampal CA1 neurons, and are consistent with NSF acting downstream of PKC ϵ in mediating downregulation of GABA $_A$ receptors.

Discussion

Our results identify a PKC ϵ –NSF signaling pathway that regulates inhibitory synapses by promoting GABA $_A$ receptor downregulation. We found that PKC ϵ phosphorylates NSF in its catalytic domain, thereby increasing NSF ATPase activity. Several upstream activators known to regulate GABA $_A$ receptor trafficking by a PKC-dependent mechanism could activate this signaling pathway. Examples include brain-derived neurotrophic factor in mouse superior colliculus (Henneberger et al., 2002) and rat hippocampal neurons (Brünig et al., 2001), serotonin in rat prefrontal cortical pyramidal neurons (Feng et al., 2001), and




Figure 7. NSF ATPase activity regulates cell surface levels of $\gamma 2$ subunits and evoked GABA $_A$ currents. **A**, HEKGR-AS-PKC ϵ cells were incubated with the DMSO vehicle as a control ($n = 13$), 1 μM TAT ($n = 11$), TAT plus 30 nM PMA ($n = 10$), 1 μM NSFi ($n = 8$), or NSFi plus PMA ($n = 7$). Biotinylated cell surface proteins were compared by Western blot analysis with anti- $\gamma 2$ subunit antibodies. NSFi increased the level of cell surface GABA $_A$ $\gamma 2$ subunits and blocked PMA-induced downregulation. $*p < 0.05$ compared with control vehicle (100%) by one-sample t tests. **B**, Hippocampal neurons were treated with buffer alone as control ($n = 18$), or with 1 μM each of scrambled (scr) $\psi\epsilon\text{RACK}$ ($n = 6$), scrambled (scr) NSFi ($n = 6$), $\psi\epsilon\text{RACK}$ ($n = 12$), NSFi ($n = 12$), or NSFi plus $\psi\epsilon\text{RACK}$ ($n = 12$). Cell surface GABA $_A$ $\gamma 2$ subunits were determined by ELISA using an antibody against extracellular domains of $\gamma 2$ subunits. The $\psi\epsilon\text{RACK}$ peptide decreased $\gamma 2$ immunoreactivity at the cell surface, whereas NSFi increased it and blocked the effect of $\psi\epsilon\text{RACK}$. $*p < 0.0003$ compared with control (100%) by one-sample t tests. **C, D**, Amplitudes of eIPSCs were averaged every 3 min and expressed relative to current measured in the first 3 min immediately after establishing whole-cell patch configuration. Peptides were administered in the patch pipette. Treatment with 5 μM $\psi\epsilon\text{RACK}$ ($n = 6$) decreased the eIPSC amplitude, whereas 2 μM NSFi ($n = 7$) increased it and prevented the effect of $\psi\epsilon\text{RACK}$ (NSFi plus $\psi\epsilon\text{RACK}$; $n = 5$). Results in **D** are mean values from the last 6 min of recording (24–30 min after break in). $*p < 0.05$ compared with NSFi plus $\psi\epsilon\text{RACK}$ by Newman–Keuls test. Error bars indicate SEM.




Figure 8. Model of the NSF D1 domain showing that S460 within the $\alpha 9$ helix (orange-red) lies in close proximity (~ 3.7 Å) to D425 and E428 on the $\alpha 7$ helix (gold). The side chains of S460, D425, E428, and T461 are shown in blue, and the oxygen atoms on the side chains are shown in red.

acetylcholine in rat superior cervical ganglion neurons (Brandon et al., 2002). Thus, PKC ϵ and NSF may be critical for downregulation of inhibitory neurotransmission induced by several neurotransmitters and growth factors.

Altered NSF-dependent regulation of GABA $_A$ receptor trafficking could contribute to several phenotypes in *Prkce* $^{-/-}$ mice. These mice show decreased anxiety-like behavior and suppression of the hypothalamic–pituitary–adrenal axis, both of which can be reversed by low doses of the GABA $_A$ receptor antagonist bicuculline (Hodge et al., 2002), indicating that GABA $_A$ receptor signaling is enhanced in *Prkce* $^{-/-}$ mice. Reduced NSF-stimulated GABA $_A$ receptor trafficking may explain the increased endogenous GABA tone in these mice and their supersensitivity to several different classes of GABA $_A$ receptor agonists (Hodge et al., 1999, 2002).

Phosphorylation of NSF by PKC ϵ not only increased NSF ATPase activity but also regulated the subcellular localization of NSF. Unlike most substrate–kinase interactions, which are transient, activation of PKC ϵ stimulated the formation of an NSF–PKC ϵ complex that translocated to the plasma membrane. PKC ϵ -induced translocation of NSF could be important for regulating GABA $_A$ receptor density at the cell surface since, when translocated to the cell membrane, NSF is better able to encounter cell surface GABA $_A$ receptors and regulate their trafficking.

Previously, it was thought that PKC does not contribute to constitutive GABA $_A$ receptor trafficking since nonselective PKC inhibitors do not alter the density of cell surface GABA $_A$ receptors (Chapell et al., 1998; Connolly et al., 1999). However, our finding that PKC ϵ inhibition increases cell surface $\gamma 2$ subunits and GABA efficacy indicates that PKC ϵ regulates constitutive receptor trafficking and that another PKC isozyme may exert an opposing effect. Inhibition of NSF also increased cell surface levels of $\gamma 2$

subunits in addition to preventing PKC ϵ -stimulated receptor downregulation. Together, these results reveal important roles for NSF and PKC ϵ in both constitutive and stimulated GABA $_A$ receptor trafficking.

NSF is comprised of three domains, an N-terminal domain and two homologous ATP binding domains, D1 and D2 (Zhao et al., 2007). The N-terminal domain interacts with soluble NSF attachment proteins (SNAPs) and soluble NSF attachment protein receptors (SNAREs). The D1 domain hydrolyzes ATP and the D2 domain mediates NSF homohexamization. To understand why phosphorylation of the D1 domain at S460 but not T461 enhances the catalytic activity of NSF, we modeled the D1 domain of NSF using the crystal structure of the homologous D2 domain (PDB number 1D2N) (Fig. 8) (Lenzen et al., 1998). Results using structure-based sequence alignment (<http://www.cgl.ucsf.edu/chimera>) showed that S460 and T461 lie in the middle of the $\alpha 9$ helix. Both residues are solvent accessible, but only S460 is in direct contact (~ 3.7 Å) with D425 and E428, two acidic residues in the $\alpha 7$ helix (Fig. 8). Phosphorylation at S460 on the $\alpha 9$ helix adds a negatively charged phosphate predicted to repel these nearby acidic residues in the $\alpha 7$ helix. This repulsion could alter

the conformation of adjacent $\alpha 6$ and $\alpha 8$ helices, which directly bind ATP, and thereby enhance the catalytic function of NSF when PKC ϵ is activated. In contrast, the side chain of T461 projects outward, away from other NSF residues. Thus, phosphorylation of T461 is unlikely to alter the conformation of NSF, but instead could regulate its interaction with other molecules such as PKC ϵ .

In contrast to our findings, Marsden et al. (2007) previously reported that NMDA receptor activation stimulates exocytosis of GABA $_A$ receptors that is dependent on calcium/calmodulin-dependent kinase II (CaMKII) and NSF. Using hippocampal neurons cultured from P0 rats, they found that NMDA increases the association of NSF with GABA $_A$ $\beta 2/3$ subunits, and that a peptide inhibitor of NSF blocks NMDA-induced increases in GABA $_A$ receptors at the cell surface. There are technical differences between their study and ours that could explain why our findings are different. These include choice of species and use of different peptides with different mechanisms of action. The peptide we used inhibits homo-hexamers of NSF and abolishes NSF ATPase activity (Morrell et al., 2005). The peptide inhibitor used by Marsden and colleagues mimics the NSF binding site of α - and β -SNAPs and is designed to prevent the binding of NSF to SNAPs rather than inhibit NSF activity (Lledo et al., 1998). Moreover, the kinases that we studied, PKC ϵ versus CaMKII, are different, which raises the intriguing possibility that these kinases determine whether NSF upregulates or downregulates synaptic GABA $_A$ receptors. *In vitro* NSF appears to be a substrate of CaMKII (Hirling and Scheller, 1996), but how this kinase regulates NSF function is not yet known.

Long-term depression of AMPA currents occurs through NMDA-stimulated removal of AMPA receptors from postsynaptic membranes (Hanley, 2008). This process requires the interaction of GluA2 subunits with the protein interacting with C-kinase-1 (PICK1). NSF binds to GluA2 subunits and acts as a disassembling chaperone using energy derived from ATP hydrolysis to disassociate AMPA receptors from PICK1, thereby inhibiting GluA2 endocytosis and stabilizing AMPA receptors at synapses (Hanley et al., 2002). As shown by our current findings, PKC ϵ -phosphorylated NSF instead promotes downregulation of GABA $_A$ receptors. Although the effects of NSF on these two types of receptors are quite different, they are additive in producing a net enhancement of excitatory neurotransmission. Thus, reducing PKC ϵ and NSF activity could prove beneficial for resetting the balance of excitation and inhibition in disorders associated with neuronal hyperexcitability. Our findings suggest that inhibitors of PKC ϵ and NSF could provide an approach for achieving this goal.

References

- Abramoff MD, Magelhaes PJ, Ram SJ (2004) Image processing with ImageJ. *Biophotonics Int* 11:36–42.
- Balduzzi R, Cupello A, Robello M (2002) Modulation of the expression of GABA(A) receptors in rat cerebellar granule cells by protein tyrosine kinases and protein kinase C. *Biochim Biophys Acta* 1564:263–270.
- Bishop AC, Kung C-Y, Shah K, Witucki L, Shokat KM, Liu Y (1999) Generation of monospecific nanomolar tyrosine kinase inhibitors via a chemical genetic approach. *J Am Chem Soc* 121:627–631.
- Brandon NJ, Jovanovic JN, Smart TG, Moss SJ (2002) Receptor for activated C kinase-1 facilitates protein kinase C-dependent phosphorylation and functional modulation of GABA $_A$ receptors with the activation of G-protein-coupled receptors. *J Neurosci* 22:6353–6361.
- Brose N, Rosenmund C (2002) Move over protein kinase C, you've got company: alternative cellular effectors of diacylglycerol and phorbol esters. *J Cell Sci* 115:4399–4411.
- Brüning I, Penschuck S, Berninger B, Benson J, Fritschy JM (2001) BDNF reduces miniature inhibitory postsynaptic currents by rapid downregulation of GABA $_A$ receptor surface expression. *Eur J Neurosci* 13:1320–1328.
- Chapell R, Bueno OF, Alvarez-Hernandez X, Robinson LC, Leidenheimer NJ (1998) Activation of protein kinase C induces gamma-aminobutyric acid type A receptor internalization in *Xenopus* oocytes. *J Biol Chem* 273:32595–32601.
- Charych EI, Liu F, Moss SJ, Brandon NJ (2009) GABA $_A$ receptors and their associated proteins: implications in the etiology and treatment of schizophrenia and related disorders. *Neuropharmacology* 57:481–495.
- Chen L, Hahn H, Wu G, Chen CH, Liron T, Schechtman D, Cavallaro G, Banci L, Guo Y, Bolli R, Dorn GW 2nd, Mochly-Rosen D (2001) Opposing cardioprotective actions and parallel hypertrophic effects of delta PKC and epsilon PKC. *Proc Natl Acad Sci U S A* 98:11114–11119.
- Chen ZW, Olsen RW (2007) GABA $_A$ receptor associated proteins: a key factor regulating GABA $_A$ receptor function. *J Neurochem* 100:279–294.
- Cinar H, Barnes EM Jr (2001) Clathrin-independent endocytosis of GABA $_A$ receptors in HEK 293 cells. *Biochemistry* 40:14030–14036.
- Collingridge GL, Isaac JT, Wang YT (2004) Receptor trafficking and synaptic plasticity. *Nat Rev Neurosci* 5:952–962.
- Connolly CN, Kittler JT, Thomas P, Uren JM, Brandon NJ, Smart TG, Moss SJ (1999) Cell surface stability of gamma-aminobutyric acid type A receptors. Dependence on protein kinase C activity and subunit composition. *J Biol Chem* 274:36565–36572.
- Essrich C, Lorez M, Benson JA, Fritschy JM, Lüscher B (1998) Postsynaptic clustering of major GABA $_A$ receptor subtypes requires the gamma 2 subunit and gephyrin. *Nat Neurosci* 1:563–571.
- Feng J, Cai X, Zhao J, Yan Z (2001) Serotonin receptors modulate GABA $_A$ receptor channels through activation of anchored protein kinase C in prefrontal cortical neurons. *J Neurosci* 21:6502–6511.
- French AP, Mills S, Swarup R, Bennett MJ, Pridmore TP (2008) Colocalization of fluorescent markers in confocal microscope images of plant cells. *Nat Protoc* 3:619–628.
- Goodkin HP, Joshi S, Mtchedlishvili Z, Brar J, Kapur J (2008) Subunit-specific trafficking of GABA $_A$ receptors during status epilepticus. *J Neurosci* 28:2527–2538.
- Goto H, Terunuma M, Kanematsu T, Misumi Y, Moss SJ, Hirata M (2005) Direct interaction of N-ethylmaleimide-sensitive factor with GABA $_A$ receptor beta subunits. *Mol Cell Neurosci* 30:197–206.
- Hanley JG (2008) PICK1: a multi-talented modulator of AMPA receptor trafficking. *Pharmacol Ther* 118:152–160.
- Hanley JG, Khatri L, Hanson PI, Ziff EB (2002) NSF ATPase and alpha/beta-SNAPs disassemble the AMPA receptor-PICK1 complex. *Neuron* 34:53–67.
- Henneberger C, Jüttner R, Rothe T, Grantyn R (2002) Postsynaptic action of BDNF on GABAergic synaptic transmission in the superficial layers of the mouse superior colliculus. *J Neurophysiol* 88:595–603.
- Herring D, Huang R, Singh M, Dillon GH, Leidenheimer NJ (2005) PKC modulation of GABA $_A$ receptor endocytosis and function is inhibited by mutation of a dileucine motif within the receptor beta 2 subunit. *Neuropharmacology* 48:181–194.
- Hirling H, Scheller RH (1996) Phosphorylation of synaptic vesicle proteins: modulation of the alpha SNAP interaction with the core complex. *Proc Natl Acad Sci U S A* 93:11945–11949.
- Hodge CW, Mehmert KK, Kelley SP, McMahon T, Haywood A, Olive MF, Wang D, Sanchez-Perez AM, Messing RO (1999) Supersensitivity to allosteric GABA $_A$ receptor modulators and alcohol in mice lacking PKC ϵ . *Nat Neurosci* 2:997–1002.
- Hodge CW, Raber J, McMahon T, Walter H, Sanchez-Perez AM, Olive MF, Mehmert K, Morrow AL, Messing RO (2002) Decreased anxiety-like behavior, reduced stress hormones, and neurosteroid supersensitivity in mice lacking protein kinase C ϵ . *J Clin Invest* 110:1003–1010.
- Kalueff AV, Nutt DJ (2007) Role of GABA in anxiety and depression. *Depress Anxiety* 24:495–517.
- Khasar SG, Lin YH, Martin A, Dadgar J, McMahon T, Wang D, Hundle B, Aley KO, Isenberg W, McCarter G, Green PG, Hodge CW, Levine JD, Messing RO (1999) A novel nociceptor signaling pathway revealed in protein kinase C epsilon mutant mice. *Neuron* 24:253–260.
- Kittler JT, Moss SJ (2003) Modulation of GABA $_A$ receptor activity by phosphorylation and receptor trafficking: implications for the efficacy of synaptic inhibition. *Curr Opin Neurobiol* 13:341–347.
- Kittler JT, Rostaing P, Schiavo G, Fritschy JM, Olsen R, Triller A, Moss SJ

- (2001) The subcellular distribution of GABARAP and its ability to interact with NSF suggest a role for this protein in the intracellular transport of GABA(A) receptors. *Mol Cell Neurosci* 18:13–25.
- Knabl J, Witschi R, Hösl K, Reinold H, Zeilhofer UB, Ahmadi S, Brockhaus J, Sergejeva M, Hess A, Brune K, Fritschy JM, Rudolph U, Möhler H, Zeilhofer HU (2008) Reversal of pathological pain through specific spinal GABA_A receptor subtypes. *Nature* 451:330–334.
- Kumar S, Fleming RL, Morrow AL (2004) Ethanol regulation of gamma-aminobutyric acid A receptors: genomic and nongenomic mechanisms. *Pharmacol Ther* 101:211–226.
- Leidenheimer NJ (2008) Regulation of excitation by GABA_A receptor internalization. *Results Probl Cell Differ* 44:1–28.
- Leidenheimer NJ, Chapell R (1997) Effects of PKC activation and receptor desensitization on neurosteroid modulation of GABA_A receptors. *Brain Res Mol Brain Res* 52:173–181.
- Leidenheimer NJ, McQuilkin SJ, Hahner LD, Whiting P, Harris RA (1992) Activation of protein kinase C selectively inhibits the gamma-aminobutyric acid A receptor: role of desensitization. *Mol Pharmacol* 41:1116–1123.
- Leil TA, Chen ZW, Chang CS, Olsen RW (2004) GABA_A receptor-associated protein traffics GABA_A receptors to the plasma membrane in neurons. *J Neurosci* 24:11429–11438.
- Lenzen CU, Steinmann D, Whiteheart SW, Weis WI (1998) Crystal structure of the hexamerization domain of N-ethylmaleimide-sensitive fusion protein. *Cell* 94:525–536.
- Lledo PM, Zhang X, Südhof TC, Malenka RC, Nicoll RA (1998) Postsynaptic membrane fusion and long-term potentiation. *Science* 279:399–403.
- Marsden KC, Beattie JB, Friedenthal J, Carroll RC (2007) NMDA receptor activation potentiates inhibitory transmission through GABA receptor-associated protein-dependent exocytosis of GABA_A receptors. *J Neurosci* 27:14326–14337.
- Matveeva EA, Whiteheart SW, Vanaman TC, Slevin JT (2001) Phosphorylation of the N-ethylmaleimide-sensitive factor is associated with depolarization-dependent neurotransmitter release from synaptosomes. *J Biol Chem* 276:12174–12181.
- Meier J, Akyeli J, Kirischuk S, Grantyn R (2003) GABA_A receptor activity and PKC control inhibitory synaptogenesis in CNS tissue slices. *Mol Cell Neurosci* 23:600–613.
- Morrell CN, Matsushita K, Lowenstein CJ (2005) A novel inhibitor of N-ethylmaleimide-sensitive factor decreases leukocyte trafficking and peritonitis. *J Pharmacol Exp Ther* 314:155–161.
- Olsen RW, Sieghart W (2008) International Union of Pharmacology. LXX. Subtypes of gamma-aminobutyric acid(A) receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* 60:243–260.
- Qi ZH, Song M, Wallace MJ, Wang D, Newton PM, McMahon T, Chou WH, Zhang C, Shokat KM, Messing RO (2007) Protein kinase C epsilon regulates gamma-aminobutyrate type A receptor sensitivity to ethanol and benzodiazepines through phosphorylation of gamma2 subunits. *J Biol Chem* 282:33052–33063.
- Qian Y, Galli A, Ramamoorthy S, Risso S, DeFelice LJ, Blakely RD (1997) Protein kinase C activation regulates human serotonin transporters in HEK-293 cells via altered cell surface expression. *J Neurosci* 17:45–57.
- Song M, Messing RO (2005) Protein kinase C regulation of GABA_A receptors. *Cell Mol Life Sci* 62:119–127.
- Violin JD, Zhang J, Tsien RY, Newton AC (2003) A genetically encoded fluorescent reporter reveals oscillatory phosphorylation by protein kinase C. *J Cell Biol* 161:899–909.
- Zhao C, Slevin JT, Whiteheart SW (2007) Cellular functions of NSF: not just SNAPs and SNAREs. *FEBS Lett* 581:2140–2149.