

Endosomal Phosphatidylinositol 3-Kinase Is Essential for Canonical GPCR Signaling[§]

Yasunori Uchida, Florentine U. Rutaganira, Damien Jullié, Kevan M. Shokat, and Mark von Zastrow

Department of Psychiatry (Y.U., D.J., and M.Z.), Department of Cellular and Molecular Pharmacology (F.U.R., K.M.S., and M.Z.), and Howard Hughes Medical Institute (F.U.R. and K.M.S.), University of California, San Francisco, San Francisco, California

Received July 20, 2016; accepted November 4, 2016

ABSTRACT

G protein-coupled receptors (GPCRs), the largest family of signaling receptors, are critically regulated by endosomal trafficking, suggesting that endosomes might provide new strategies for manipulating GPCR signaling. Here we test this hypothesis by focusing on class III phosphatidylinositol 3-kinase (Vps34), which is an essential regulator of endosomal trafficking. We verify that Vps34 is required for recycling of the β 2-adrenoceptor (β 2AR), a prototypical GPCR, and then investigate the effects of Vps34 inhibition on the canonical cAMP response elicited by β 2AR

activation. Vps34 inhibition impairs the ability of cells to recover this response after prolonged activation, which is in accord with the established role of recycling in GPCR resensitization. In addition, Vps34 inhibition also attenuates the short-term cAMP response, and its effect begins several minutes after initial agonist application. These results establish Vps34 as an essential determinant of both short-term and long-term canonical GPCR signaling, and support the potential utility of the endosomal system as a druggable target for signaling.

Introduction

Many signaling receptors, including G protein-coupled receptors (GPCRs), enter the endosomal system after ligand-induced activation at the plasma membrane (PM), and the endosomal network is known to have myriad effects on cellular GPCR signaling (Pierce et al., 2002; Sorkin and von Zastrow, 2009; von Zastrow and Williams, 2012; Irannejad et al., 2015). Internalized receptors are sorted after endocytosis for recycling back to the PM or subsequent transport to lysosomes for degradation. This sorting decision, by determining whether regulated endocytosis maintains or depletes the surface receptor complement, has been believed to exert long-term control over cellular signaling responsiveness under conditions of prolonged or repeated ligand application (Hanyaloglu and von Zastrow, 2008). In addition, endosomes themselves can function as sites of receptor-mediated signal initiation,

which potentially affects both short-term and long-term cellular signaling activities (Murphy et al., 2009; Irannejad and von Zastrow, 2014).

Phosphatidylinositol (PI) 3-phosphate (PI3P) is a phosphorylated inositol lipid (phosphoinositide) that is found mainly in endosomal membranes (Gillooly et al., 2000). This lipid, although present in relatively small amounts, is a defining feature of endosomal membranes that regulates many aspects of endosome biogenesis, organization, and trafficking by recruiting PI3P-binding proteins (Schink et al., 2013). A major pathway of PI3P synthesis is mediated by class III phosphatidylinositol 3-kinase (Vps34), which specifically phosphorylates PI to produce PI3P in endosomal membranes (Shin et al., 2005; Raiborg et al., 2013).

β 2-adrenoceptors (β 2ARs) are widely considered to be prototypic GPCRs (Lefkowitz, 2007). β 2-adrenoceptors internalize rapidly after ligand-induced activation via clathrin-coated pits and have the ability to recycle to the PM with remarkably high efficiency via a retromer-dependent pathway (Goodman et al., 1996; Temkin et al., 2011). Several components of this machinery require 3-phosphorylated phosphoinositides for membrane attachment (Cullen and Korswagen, 2011), and a previous study (Awwad et al., 2007) has implicated Vps34 in promoting β 2AR recycling. Accordingly, we hypothesized that

This work was supported by the National Institutes of Health [Grants DA-010711 and DA-012864, R01-AI-099245, and U19-AI-109622]. Y.U. is the recipient of a Postdoctoral Fellowship for Research Abroad from Japan Society for the Promotion of Science and a Research Fellowship from the Uehara Memorial Foundation.

dx.doi.org/10.1124/mol.116.106252.

[§] This article has supplemental material available at molpharm.aspetjournals.org.

ABBREVIATIONS: ANOVA, analysis of variance; AR, adrenoceptor; β 2ARs, β 2-adrenoceptors; DMEM, Dulbecco's modified Eagle's medium; DMSO, dimethylsulfoxide; FKBP, FK506 binding protein, FRB, FKBP12-rapamycin-binding domain; GDC-0941, 4-[2-(1H-indazol-4-yl)-6-[(4-methylsulfonyl)piperazin-1-yl)methyl]thieno[3,2-d]pyrimidin-4-yl]morpholine; GFP, green fluorescent protein; GPCR, G protein-coupled receptor; HEK, human embryonic kidney; PBS, phosphate-buffered saline; PI, phosphatidylinositol; PI3K, phosphatidylinositol 3-kinase; PI3P, phosphatidylinositol 3-phosphate; PI(3,5)P₂, phosphatidylinositol 3,5-bisphosphate; PM, plasma membrane; SAR405, (8S)-9-[(5-chloropyridin-3-yl)methyl]-2-[(3R)-3-methylmorpholin-4-yl]-8-(trifluoromethyl)-7,8-dihydro-6H-pyrimido[1,2-a]pyrimidin-4-one; UCSF, University of California, San Francisco; Vps34, class III phosphatidylinositol 3-kinase; VPS34-IN1, 1-((2-((2-chloropyridin-4-yl)amino)-4'-(cyclopropylmethyl)-[4,5'-bipyrimidin]-2'-yl)amino)-2-methylpropan-2-ol; WM, wortmannin; WT, wild type; YM201636, 6-amino-N-[3-(4-morpholin-4-yl)pyrido[2,3-furo[2,4-b]pyrimidin-2-yl)phenyl]pyridine-3-carboxamide.

Vps34 represents an endosomal target that controls β 2AR signaling. Chemical inhibitors used in previous studies of β 2AR trafficking (Sorensen et al., 1999; Naga Prasad et al., 2001; Awwad et al., 2007) [wortmannin (WM) and LY294002] have additional cellular targets (Knight, 2010). In the current study, we employed more specific manipulations to examine the trafficking and signaling effects of endosomal PI3P and Vps34, and revealed previously unrecognized roles of this critical endosomal kinase in GPCR signaling. These results demonstrate that both short-term and sustained GPCR signaling can be manipulated by targeting the endosomal system.

Materials and Methods

Cell Culture, cDNA Constructs, and Transfection. Human embryonic kidney (HEK) 293 cells (American Type Culture Collection, Manassas, VA) were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum [University of California, San Francisco (UCSF) Cell Culture Facility, San Francisco, CA]. A plasmid encoding Flag-tagged β 2AR was previously described (Cao et al., 1999). A plasmid encoding enhanced green fluorescent protein (GFP)-tagged Hrs 2xFYVE was a gift from Harald Stenmark (Oslo University Hospital, Oslo, Norway) (Gillooly et al., 2000). mCherry-FKBP-myotubularin 1 (MTM1) [wild type (WT)] and iRFP-FRB (FKBP12-rapamycin-binding domain)-Rab5 plasmids were obtained from Tamas Balla (National Institutes of Health, Bethesda, MD) through Addgene (Cambridge, MA) (Hammond et al., 2014). A phosphatase-dead mutant of mCherry-FKBP-MTM1 (C375S) was generated by using Quikchange Site-Directed Mutagenesis Kit (Agilent, Santa Clara, CA). Nb37-GFP plasmid was previously described (Irannejad et al., 2013). DNA transfection was performed by using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) according to the manufacturer instructions, and cells were used for subsequent experiments 24 hours after transfection, unless otherwise indicated. Cells stably expressing with Flag-tagged β 2AR were created as previously described (Laufer et al., 2010).

Inhibitors. PIK-III was synthesized according to the published protocol (Honda et al., 2015). The commercial sources of other inhibitors used in this study were as follows: WM (Sigma-Aldrich, St. Louis, MO); VPS34-IN1 (1-((2-(2-chloropyridin-4-yl)amino)-4'-(cyclopropylmethyl)-[4,5'-bipyrimidin]-2'-yl)amino)-2-methylpropan-2-ol; MedKoo Biosciences, Chapel Hill, NC); SAR405 ((8S)-9-[5-chloropyridin-3-yl)methyl]-2-[(3R)-3-methylmorpholin-4-yl]-8-(trifluoromethyl)-7,8-dihydro-6H-pyrimido[1,2-a]pyrimidin-4-one; ApexBio, Houston, TX); GDC-0941 (4-[2-(1H-indazol-4-yl)-6-[4-methylsulfonylpiperazin-1-yl)methyl]thieno[3,2-d]pyrimidin-4-yl]morpholine; Selleck Chemicals, Houston, TX); and YM201636 (YM201636, 6-amino-N-[3-(4-morpholin-4-yl)pyrido[2,3-furo[2,4-b]pyrimidin-2-yl)phenyl]pyridine-3-carboxamide; Cayman Chemical, Ann Arbor, MI). Unless otherwise indicated, cells were pretreated with these inhibitors in serum-free DMEM for either 1 hour (100 nM WM, 3 μ M PIK-III, 1 μ M VPS34-IN1, 3 μ M SAR405, or 1 μ M GDC-0941) or 3 hours (800 nM YM201636) before agonist application. Trafficking and signaling assays were performed in the presence of the indicated inhibitors.

Trafficking Assays. Unless otherwise indicated, trafficking assays were performed at 37°C using HEK293 cells stably expressing Flag- β 2AR. Cells plated on coverslips were used to assess receptor localization by fluorescence microscopy. To examine agonist-induced receptor internalization, cells were treated with 10 μ M isoproterenol (a β 2AR agonist; Sigma-Aldrich) for 25 minutes. To examine recycling after agonist removal, cells were first treated with isoproterenol for 25 minutes and then washed with phosphate-buffered saline (PBS) and treated with 10 μ M alprenolol (a β 2AR antagonist; Sigma-Aldrich) for 45 minutes. Antagonist was used to prevent any residual agonist effects in the recycling period. Cells were fixed by 4% paraformaldehyde in PBS for 20 minutes, quenched with

Tris-buffered saline for 20 minutes, and subjected to immunocytochemistry, as described below. For acute PI3P depletion by MTM1 recruitment, cells transfected with mCherry-FKBP-MTM1 and iRFP-FRB-Rab5 plasmids were used, and 1 μ M rapamycin (Sigma-Aldrich) was applied to cells either 20 minutes before isoproterenol treatment (for internalization) or with the same timing as for alprenolol (for recycling).

Cells were plated on 12-well plates to analyze β 2AR trafficking by fluorescence flow cytometry. Both agonist-induced internalization and recycling after agonist removal were tracked quantitatively by determining surface β 2AR levels. For internalization, cells were treated with 1 μ M isoproterenol for the indicated times. For recycling, cells were first treated with 1 μ M isoproterenol for 25 minutes and then with 10 μ M alprenolol for 45 minutes. Surface β 2AR receptors were then labeled with Alexa Fluor 488-conjugated M1 anti-Flag antibody (Sigma-Aldrich) at 4°C. The mean fluorescence intensity of each sample (2000–10,000 cells/sample) was measured using a FACSCalibur flow cytometer (BD Biosciences, San Jose, CA). The percentage of recycled receptors was calculated as follows: $100 \times [(intensity\ after\ the\ recycling\ period) - (intensity\ after\ the\ internalization\ period)] / [(intensity\ without\ agonist\ addition) - (intensity\ after\ the\ internalization\ period)]$.

To measure β 2AR recycling in the continuous presence of agonist, a previously established method (Tsao and von Zastrow, 2000) was used with modifications. Briefly, surface β 2ARs were first labeled for 10 minutes with Alexa Fluor 647-conjugated M1 anti-Flag antibody, which requires calcium to bind the Flag epitope. Cells were then treated with 1 μ M isoproterenol for 25 minutes to induce receptor internalization. At this point, cells were washed with calcium, and magnesium-free PBS with 0.4% EDTA to dissociate antibodies from surface receptors and specifically label internalized receptors. Cells were then incubated in EDTA-supplemented PBS with 1 μ M isoproterenol for 5 minutes. Control samples without this incubation period were also made. Cells were then chilled on ice and subjected to flow cytometry, as described above, to measure the antibody efflux in the 5-minute period. Because antibodies bound to receptors that were recycled to the PM in this period eluted immediately, the antibody efflux was correlated to the degree of β 2AR recycling.

Immunocytochemistry and Fixed-Cell Imaging. Permeabilization and blocking were performed for 20 minutes with 0.1% Triton X-100 and 3% skimmed milk in PBS. Cells were then incubated with anti-Flag (M1) antibody (1:1000; Sigma-Aldrich) and Alexa Fluor 488-conjugated secondary antibody (1:1000; Molecular Probes, Carlsbad, CA) in the blocking buffer for 1 hour each. Fixed samples were mounted with ProLong Gold (Molecular Probes). Cells were imaged with an inverted microscope (TE-2000; Nikon, Tokyo, Japan) with a confocal scanner unit (CSU22; Yokogawa, Tokyo, Japan) using a 100 \times numerical aperture 1.45 objective. Images were collected using an electron-multiplying CCD camera (iXon 897; Andor Technology, Belfast, UK) operated in the linear range controlled by Micro-Manager software (<https://www.micro-manager.org>).

Live-Cell Confocal Imaging. Cells were imaged in DMEM without phenol red (UCSF Cell Culture Facility) with 30 mM HEPES (pH 7.4). For MTM1 recruitment experiments, HEK293 cells stably expressing Flag- β 2AR were used and imaged with the aforementioned confocal microscope. For Nb37 localization experiments, HEK293 cells were transiently transfected with Flag- β 2AR and Nb37-enhanced GFP plasmids. Cells were then pretreated with dimethylsulfoxide (DMSO) or PIK-III, and then treated with 10 μ M isoproterenol. Receptor-expressing cells were randomly chosen for assessing Nb37 localization and imaged after 15–30 minutes of isoproterenol treatment. Cells were imaged with another spinning disk confocal microscope (Ti-E microscope; Nikon) in the Nikon Imaging Center at UCSF (with CSU22 confocal scanner unit; Yokogawa) using a 100 \times numerical aperture 1.49 objective. Images were collected using an electron-multiplying CCD camera (Evolve Delta; Photometrics, Tucson, AZ) operated in the linear range controlled by Micro-Manager software.

Image Analysis. Images were saved as 16-bit TIFF files and analyzed using Fiji software (Schindelin et al., 2012). Colocalization between Flag- β 2AR and FRB-Rab5 was estimated by calculating Pearson's correlation coefficients between the two channels using the Coloc 2 plug-in in Fiji. Line-scan analysis was performed with the plot profile function, and the obtained values were normalized to the maximum value of each channel. The intensities of Nb37 and β 2AR were measured by a custom-written program created by one of the authors (D.J.), which works on MATLAB (MathWorks, Natick, MA). The script is shown in the Supplemental Material. Briefly, after selecting a background region in cytosol and β 2AR-containing endosomes, the program created donut-shaped regions (3 pixels wide), which include endosome-limiting membranes, and calculated the average fluorescence intensity of each donut. Five endosomes per cell were randomly chosen for the analysis, and endosomes with negative Nb37 intensities were excluded from the analysis. The Nb37 intensity of each endosome was then normalized to the β 2AR intensity.

Luminescence-Based Real-Time cAMP Assay. The experimental procedure was performed as described previously, except for using WT HEK293 cells (Irannejad et al., 2013). Briefly, cells were transiently transfected with pGloSensor-20F (Promega, Madison, WI), which encodes a cyclic-permuted luciferase cAMP reporter construct. For short-term signaling, cells were treated with luciferin (GoldBio, St. Louis, MO) in serum-free media for 1 hour in a 24-well

plate, and luminescence values in the experimental wells were obtained after adding 1 μ M isoproterenol. Reference wells were made in the same columns in the 24-well plate as the experimental wells and were treated with 5 μ M forskolin (Sigma-Aldrich). The luminescence values obtained in experimental wells were normalized to the maximum and minimum values in the reference wells (i.e., the maximum and minimum values of the reference wells were set to 100% and 0%, respectively). For repeated signaling, cells were lifted and seeded on a 12-well plate 1 d after transfection, and the experiments were performed on the next day. Cells were treated with DMSO or 5 μ M PIK-III and luciferin in serum-containing media for 1 hour, and then treated with 1 μ M isoproterenol for 1 hour (first treatment). Cells were washed three times with serum-containing media without isoproterenol, and 10 minutes after washout were re-challenged with 1 μ M isoproterenol (second treatment). Luminescence was recorded after both the first and second treatment, and the increase in luminescence values after the second treatment was normalized to that after the first treatment in the same well.

Biochemical cAMP Assay. After the indicated time of isoproterenol or forskolin treatment, cells were washed with ice-cold PBS and lysed by 0.1 M HCl for 10 minutes at room temperature. The cAMP concentration in the lysate was determined by using the Direct cAMP ELISA kit (Enzo Life Sciences, Farmingdale, NY) according to the manufacturer instructions. The cAMP concentration was normalized




Fig. 1. Endosomal PI3P is required for β 2AR recycling, but not for endocytosis. (A) Schematic of the rapamycin (rapa)-induced recruitment of MTM1 to Rab5-containing endosomes. (B) Representative images of live cells expressing GFP-2xFYVE (a PI3P probe protein), mCherry-FKBP-MTM1 [either WT or a phosphatase-dead mutant (C375S)] and iRFP-FRB-Rab5. Cells were imaged before and after 5 minutes of rapa treatment by confocal microscopy. Scale bar, 10 μ m. (C and D) Trafficking of β 2AR examined by immunofluorescence microscopy. (C) Cells stably expressing Flag- β 2AR were transfected with mCherry-FKBP-MTM1 (WT or C375S) and iRFP-FRB-Rab5. In the agonist condition, cells were treated with rapa for 20 minutes, and then with a β 2AR agonist isoproterenol (iso) for 25 minutes. In the agonist-to-antagonist condition, cells were first treated with isoproterenol for 25 minutes, and then with a β 2AR antagonist alprenolol and rapa for 45 minutes. After fixation, cells were stained for Flag epitope. Representative confocal images are shown. Insets show the boxed areas at higher magnification. Arrows show β 2AR localization to Rab5-containing endosomes. Scale bar, 10 μ m. (D) Quantification of the results. Pearson's correlation coefficients between β 2AR and FRB-Rab5 were calculated ($n = 11$ [WT, agonist], 11 [C375S, agonist], 21 [WT, agonist to antagonist], or 19 [C375S, agonist to antagonist] cells from two independent experiments, *** $P < 0.001$ by two-tailed t test). Error bars represent the S.D.




Fig. 2. Vps34 is required for efficient β 2AR recycling, but not for endocytosis. (A) Representative images of HEK293 cells expressing GFP-2xFYVE. Cells were treated with PIK-III or DMSO for 1 hour, fixed, and imaged by confocal microscopy. Scale bar, 10 μ m. (B) Effects of PI3K inhibitors on basal surface β 2AR levels. Cells stably expressing Flag-tagged β 2AR were treated with the indicated inhibitors for 1 hour. Surface receptor levels were then determined by flow cytometry and were expressed as a percentage of the levels in DMSO-treated cells ($n = 4$ independent experiments). (C) Effects of PI3K inhibitors on agonist-induced reduction of surface β 2AR levels. After 1 hour of pretreatment of inhibitors, cells were treated with isoproterenol for the indicated periods, and surface receptor levels were determined. Basal receptor levels shown in (B) were included in a graph as “0 minutes” for comparison, and data were expressed as a percentage of the basal receptor levels in DMSO-treated cells. Data are from $n = 3$ (5–40 minutes) or $n = 4$ (0 minutes) independent experiments. PIK-III or WM treatment significantly reduced surface β 2AR levels compared with DMSO control at 10, 20, and 40 minutes after isoproterenol treatment ($*P < 0.05$, $**P < 0.01$, $***P < 0.001$ by two-way ANOVA and Tukey’s post hoc tests). (D) The efficiency of β 2AR endocytosis was estimated from the percentage reduction in surface receptor levels after the shortest (5 minutes) isoproterenol application in (C). Data were from $n = 3$ independent experiments. (E) Direct measurement of β 2AR recycling after agonist removal and antagonist treatment. After 1 hour of treatment of the indicated inhibitor, cells were treated with isoproterenol for 25 minutes, and then with alprenolol for 45 minutes. The percentage of recycled receptors was calculated as described in *Materials and Methods* and is shown in bar graphs ($n = 3$ independent experiments, $***P < 0.001$ compared with DMSO control by one-way ANOVA and Tukey’s post hoc tests). (F) Recycling defect in PIK-III–treated cells was

to the protein concentration determined by Coomassie Plus Protein Assay (Pierce, Waltham, MA) and was displayed as the percentage of the cAMP level in the reference sample for each independent experiment. The reference sample of each experiment is described in the figure legends.

Statistics and Reproducibility. Statistical analysis was performed using Prism 6 software (GraphPad Software, La Jolla, CA). Statistical significance was determined with either two-tailed Student’s *t* test or one-way analysis of variance (ANOVA) and considered to be significant if *P* values were less than 0.05 (for details, see figure legends). Independent experiments were performed on different days. Representative microscopic data were from at least two independent experiments.

Results

To specifically investigate the effects of endosomal PI3P, we employed a system to inducibly hydrolyze endosomal PI3P through rapamycin-induced recruitment of the inositol 3-phosphatase MTM1 to endosomes in cells expressing MTM1 fused to the FK506 binding protein (FKBP) and the FRB domain fused to Rab5 (Hammond et al., 2014) (Fig. 1A). FKBP-MTM1 mainly localized in the cytoplasm but was recruited in the short term to endosomes after rapamycin addition (Fig. 1B, right panels). We confirmed effective depletion of endosomal PI3P in HEK293 cells by using a PI3P probe protein (2xFYVE domain of Hrs) (Gillooly et al., 2000). Before rapamycin addition, 2xFYVE showed punctate localization corresponding to PI3P-containing endosomal membranes. After 5 minutes of rapamycin treatment, the 2xFYVE probe lost its punctate localization and showed even distribution in cytoplasm, concomitant with FKBP-MTM1 accumulation on endosomes (Fig. 1B, left panels). This effect was dependent on the phosphatase activity of MTM1 because a phosphatase-dead mutant of MTM1 (C375S) (Fili et al., 2006) did not cause dissociation of the 2xFYVE probe after rapamycin-induced recruitment (Fig. 1B, compare WT to C375S).

Endosomal recruitment of MTM1 (WT) did not detectably perturb agonist (isoproterenol)-induced endocytosis of β 2ARs, as indicated by the robust redistribution of Flag- β 2ARs from PM to endosomes, which was indistinguishable from that observed after the recruitment of the control, phosphatase-dead MTM1 (C375S) (Fig. 1C, agonist condition, top rows). However, MTM1 (WT) recruitment impaired β 2AR recycling, which is indicated by the retention of Flag- β 2AR in endosomes after isoproterenol removal, whereas efficient recycling was observed after recruitment of the phosphatase-dead MTM1 (C375S) (Fig. 1C, agonist-to-antagonist condition, bottom panels). These results, which were verified across multiple

verified by fluorescence microscopy. After 1 hour treatment with PIK-III or DMSO, cells were treated with isoproterenol for 25 minutes (in the agonist condition) and then with alprenolol for 45 minutes (in the agonist-to-antagonist condition). After fixation, cells were stained for the Flag epitope. Representative confocal images are shown. Scale bars, 10 μ m. (G) β 2AR recycling in the continuous presence of agonist. Surface receptors were labeled with Alexa Fluor 647–conjugated anti-Flag antibody, and then internalized by 1 μ M isoproterenol for 25 minutes. After stripping antibodies bound to residual surface receptors by calcium-depleted medium, cells were further incubated for 5 minutes in calcium-depleted medium. β 2AR recycling was then estimated by measuring the antibody efflux in this 5-minute period ($n = 3$ samples from one experiment, $**P < 0.01$ by two-tailed *t* test. Similar results were obtained in two other independent experiments). Error bars indicate the S.D.

samples (Fig. 1D), suggest that endosomal PI3P is specifically required for β 2AR recycling but not endocytosis.

To examine the role of Vps34 activity, we used PIK-III, a potent chemical inhibitor of Vps34 that has high selectivity for Vps34 over other kinases (Dowdle et al., 2014; Honda et al., 2015). PIK-III application produced a pronounced redistribution of the 2xFYVE probe from endosomes to the cytoplasm (Fig. 2A). We then quantified the effects of specific Vps34 inhibition on β 2AR trafficking using fluorescence flow cytometry. PIK-III did not detectably change basal surface β 2AR levels over the time interval required for our experiments, similar to the broad-spectrum PI 3-kinase (PI3K) inhibitor WM (Fig. 2B). These inhibitors also had little effect on isoproterenol-induced reduction of β 2ARs from the PM measured 5 minutes after agonist application (Fig. 2, C and D), a time point at which net β 2AR internalization is dominated by endocytic rate. However, both PIK-III and WM significantly reduced surface concentration of β 2ARs at later time points (Fig. 2C), suggesting a selective inhibition of receptor recycling. We verified this by direct measurement of β 2AR recycling after agonist removal (Fig. 2E). The similar degree of inhibition produced by PIK-III and WM suggests that Vps34 is the major PI3K isoform controlling β 2AR trafficking in this cell system. We also confirmed the PIK-III effects by immunofluorescence microscopy. PIK-III caused clearly detectable retention of β 2AR in intracellular structures after agonist removal (Fig. 2F), consistent with the recycling defect observed by flow cytometry.

Internalized β 2ARs are efficiently recycled not only after agonist removal, but also in the continuous presence of agonist (Morrison et al., 1996; Tsao and von Zastrow, 2000). To analyze β 2AR recycling in the presence of agonist, we employed a previously established “loss of internal receptor” assay (Tsao and von Zastrow, 2000). In this assay, Flag-tagged β 2ARs are labeled with fluorochrome-conjugated anti-Flag antibody and are subsequently internalized by isoproterenol application. The recycling of Flag-tagged β 2AR is then estimated by the efflux of anti-Flag antibody to calcium-depleted medium. PIK-III essentially abolished this antibody efflux, providing evidence that PIK-III blocks β 2AR recycling in the presence of isoproterenol (Fig. 2G).




Fig. 3. Vps34, but not class I PI3Ks or PIKfyve, is required for β 2AR recycling. After 1 hour (A) or 3 hours (B) of treatment with the indicated inhibitor, cells were subjected to the recycling assay as in Fig. 2E [$n = 7$ (DMSO)], or $n = 3$ (each inhibitor) independent experiments in (A), and $n = 3$ independent experiments in (B). *** $P < 0.001$ compared with DMSO control by one-way ANOVA and Tukey's post hoc tests). Error bars indicate the S.D.

To further confirm the role of Vps34 in β 2AR recycling, we used two additional recently reported selective Vps34 inhibitors, VPS34-IN1 (Bago et al., 2014) and SAR405 (Ronan et al., 2014). Both inhibitors suppressed β 2AR recycling after agonist removal as well as PIK-III (Fig. 3A). In marked contrast, a selective inhibitor of class I PI3Ks, GDC-0941 (Folkes et al., 2008), did not affect β 2AR recycling, indicating that class I PI3Ks are not required for β 2AR recycling (Fig. 3A).

Notably, PI3P is a precursor for producing another phosphoinositide phosphatidylinositol 3,5-bisphosphate (PI(3,5)P₂), which also regulates trafficking and ion homeostasis at endosomes (McCartney et al., 2014). The reduction of PI3P levels thus potentially affects PI(3,5)P₂ production, and, moreover, MTM1 has a phosphatase activity toward PI(3,5)P₂ as well as PI3P (Fili et al., 2006). Thus, we considered the possibility that Vps34 inhibition affects β 2AR recycling through depleting PI(3,5)P₂ rather than PI3P itself. To test this, we used YM201636 to selectively block PIKfyve, which generates PI(3,5)P₂ from PI3P (Jefferies et al., 2008), and measured β 2AR recycling by flow cytometry. YM201636, in contrast to PIK-III, did not significantly affect β 2AR recycling (Fig. 3B). These observations verify that PI3P is the key phosphoinositide product mediating the Vps34 effect on β 2AR recycling.

We next investigated whether Vps34 affects β 2AR signaling. The key biochemical step initiating canonical β 2AR signaling is receptor-mediated activation of the heterotrimeric G protein G_s (Rasmussen et al., 2011). Ligand-activated β 2ARs activate G_s first in the PM and then in the endosome-limiting membrane after endocytosis, with both events occurring sequentially over a period of several minutes (Irannejad et al., 2013). Accordingly, we reasoned that Vps34 might affect canonical β 2AR signaling by enhancing or inhibiting the endosome phase of G_s activation. To test this, we applied a previously described conformational biosensor, Nb37-GFP, which detects G_s activation in intact cells by local recruitment (Irannejad et al., 2013). The recruitment of Nb37-GFP, which is indicative of G_s activation, was clearly evident at β 2AR-containing endosomes (Supplemental Fig. 1A) and was localized to the limiting membrane (Supplemental Fig. 1A, insets, line scans). PIK-III did not detectably change this behavior (Supplemental Fig. 1A). We quantified the intensity of Nb37-GFP signal in endosome-limiting membranes and revealed that Nb37-GFP was similarly recruited, irrespective of PIK-III application (Supplemental Fig. 1B). This suggests that Vps34 is not essential for G_s activation by β 2AR at endosomes.

After β 2AR activation, an initial phase of cytoplasmic cAMP accumulation occurs from the PM and then receptors internalize to initiate a second signaling phase from endosomes (Irannejad et al., 2013; Tsvetanova and von Zastrow, 2014). The cellular cAMP response is desensitized by prolonged agonist exposure (Pierce et al., 2002), and recycling of internalized β 2ARs to the PM after agonist removal promotes resensitization of the cellular cAMP response to subsequent agonist challenge (Yu et al., 1993; Pippig et al., 1995; Odley et al., 2004). Accordingly, to investigate the potential signaling consequences of Vps34 activity, we investigated the effects of specific Vps34 blockade, and used three different experimental protocols to examine both short-term and long-term β 2AR-mediated cAMP responses.

We first examined the isoproterenol-induced cAMP response elicited within seconds after short-term application, which we




Fig. 4. Vps34 supports not only long-term cellular β 2AR signaling but also the short-term signaling response. (A) Luminescence-based cAMP assay examining the initial accumulation of cytoplasmic cAMP induced by isoproterenol (iso) application. WT HEK293 cells were transiently transfected with pGloSensor-20F. Cells were pretreated with PIK-III or DMSO, and then iso or forskolin (fsk) was added (time = 0 seconds). cAMP responses in iso-treated cells are shown. Data are normalized to the luminescence of fsk-treated wells ($n = 6$ independent experiments). (B) Luminescence-based cAMP assay examining the resensitization of the cellular cAMP response after prolonged iso treatment and washout. Cells were pretreated with PIK-III or DMSO, and then treated with iso for 1 hour. After a 10-minute washout period, cells were rechallenged with iso. Luminescence increased by the second stimulation was normalized to that of the first stimulation. Normalized luminescence is significantly reduced in PIK-III-treated cells after 70–150 seconds of iso application compared with DMSO-treated cells ($n = 3$ independent experiments; $*P < 0.05$ by two-tailed t test). (C and D) Biochemical determination of cAMP levels. HEK293 cells were pretreated with the indicated inhibitor, and then treated with iso (C) or fsk (D) for the indicated times. Cells were then lysed, and cAMP concentration in the lysates was determined by enzyme-linked immunosorbent assay and normalized to protein concentration. In (C), data are expressed as a percentage of the cAMP levels in cells pretreated with DMSO and then with iso for 1 minute ($n = 8$ [DMSO] or $n = 4$ [PIK-III and VPS34-IN1] independent experiments; $*P < 0.05$, $**P < 0.01$ compared with DMSO control by one-way ANOVA and Tukey's post hoc tests). In (D), data are expressed as a percentage of the cAMP levels in cells pretreated with DMSO and then fsk for the indicated times ($n = 4$ independent experiments). Error bars indicate the S.E.M.

reasoned would primarily sample signaling from the PM. To do so, we used a biosensor (GloSensor; Promega) that offers rapid detection without cell lysis for sample preparation, as is required for standard biochemical determination of cAMP (Fan et al., 2008). Blocking Vps34 activity with PIK-III did not detectably affect this immediate signal (Fig. 4A). We next examined the isoproterenol response elicited after a 60-minute interval of isoproterenol pre-exposure followed by washout, which we reasoned would sample the resensitized β 2AR pool present in the PM. Vps34 blockade significantly reduced this response (Fig. 4B). Together, these results are consistent with an essential role of Vps34 in β 2AR recycling and the previously established role of recycling in promoting β 2AR resensitization.

We then investigated the effect of Vps34 blockade on the short-term isoproterenol-induced cAMP response when measured 5–10 minutes after initial application, which we reasoned could include signaling initiated from both the PM and endosomes. It was possible to use a standard biochemical determination of cAMP over this time scale, facilitating a more direct and quantitative comparison of Vps34 effects on the cAMP response than using the GloSensor. PIK-III had no detectable effect on the cAMP response measured 1 minute after isoproterenol application, which is consistent with the GloSensor results. However, when measured after 5–10 minutes, PIK-III caused a pronounced inhibition (Fig. 4C). Another selective Vps34 inhibitor, VPS34-IN1, caused essentially the same effects as PIK-III

Discussion

Since it was first identified in yeast as a gene required for vacuolar protein sorting (Herman and Emr, 1990), Vps34 has long been recognized to regulate endosomal trafficking. In line with this, a previous study (Awwad et al., 2007) showed that broad-spectrum PI3K inhibitors (LY294002 and WM), which block Vps34 and other PI3Ks, suppress β 2AR recycling. However, since such inhibitors are not specific for Vps34, it was not possible previously to interrogate the role of Vps34 or endosomal PI3P specifically. In this study, we verify the role of Vps34 and PI3P by using recently developed specific manipulations. We demonstrate that both short-term PI3P depletion in endosomal membranes (Fig. 1) and specific Vps34 inhibition (Figs. 2 and 3) suppress β 2AR recycling. In addition, we show that a highly selective inhibitor of class I PI3Ks, which are blocked by broad-spectrum inhibitors, does not affect β 2AR recycling (Fig. 3A). Together, our results unambiguously show that Vps34 is essential for β 2AR recycling. Although Vps34 might have additional kinase-independent effects, our kinase inhibitor and short-term PI3P depletion data that are presented here indicate that the kinase activity of Vps34 and endosomal PI3P are required for β 2AR recycling.

We then demonstrate a fundamental role of Vps34 in the cAMP signaling response triggered by β 2AR activation. Figure 5 summarizes our proposed model regarding this. The short-term cAMP response by single agonist application reaches a maximum before 5 minutes at the latest, but then is rapidly desensitized. After agonist washout, cellular responsiveness is recovered at least in part by β 2AR recycling to the PM. Vps34, by promoting β 2AR recycling, supports this resensitization (Fig. 5A). Our data also reveal an additional role of Vps34 in promoting the short-term cellular cAMP response, and they resolve a second component of the short-term response that is specifically Vps34 dependent (Fig. 5B).

What is the underlying mechanism of this Vps34-dependent component? We previously showed that β 2AR recycling events are observed after 3–5 minutes of agonist application, and reach steady state within 10 minutes (Yudowski et al., 2009). This is consistent with the time course of the Vps34-dependent component of short-term signaling that is established in the present work. Also, our data indicate that Vps34 is required for β 2AR recycling in the continuous presence of agonist (Fig. 2G). We thus speculate that VPS34/PI3P-mediated fast recycling enhances cAMP production (Fig. 5C). β 2AR recycling supplies receptors to the PM, and the recycled receptors are subjected to a further round of endocytosis. This rapid cycling between two compartments would sustain cAMP responses by supplying functional receptors to the PM and then to endosomes. Alternatively, although Vps34 inhibition does not detectably affect endosomal G_s activation (Supplemental Fig. 1), it could affect endosomal cAMP production downstream of G_s activation. Indeed, after 5 minutes of isoproterenol application, Vps34 inhibition had little effect on the surface β 2AR density (Fig. 2C) but reduced the cAMP levels by ~40% (Fig. 4C). Thus, it is possible that Vps34 inhibition reduces the short-term cAMP response separately from receptor recycling at this time point, presumably by affecting the endosome signal (Fig. 5C). Further studies will be needed to clarify precisely how Vps34 impacts the short-term cAMP response.




Fig. 5. Summary and model of Vps34 effects in β 2AR-mediated cAMP response. (A) There are two aspects of cAMP response that are triggered by the application of the β 2AR agonist isoproterenol (iso): one is the short-term response by single application, and the other is repeated response by a second (or more) application (boxed by a dotted line) after initial application and washout. The time course of agonist application and washout is indicated by black/white boxes above the graphs. In both cases, cAMP responses reach maxima in less than 5 minutes and are rapidly desensitized. β 2AR recycling promotes resensitization of the cellular responsiveness to subsequent agonist application in the washout period, and thus supports the repeated response. We demonstrated that efficient β 2AR recycling requires Vps34, and, in line with this, Vps34 supports the repeated cAMP response. (B and C) According to our data showing Vps34 inhibition reduces cAMP levels after 5–10 minutes of short-term agonist application, the short-term cAMP response seems to be divided into the following two phases: the Vps34-independent initial phase and the Vps34-dependent later phase. Our models regarding the role of Vps34 in the short-term cAMP response are shown in (C). In the initial phase, β 2ARs activate cAMP synthesis in the PM and endosomes, but are not recycled yet. In the later phase, the recycling of β 2AR begins in a Vps34/PI3P-dependent manner, and the recycled receptors are subjected to the next round of receptor cycling between the PM and endosomes (indicated by red arrows). The continuous cycling of β 2ARs, which is driven by Vps34, would enhance cAMP production from both locations in the later phase. It is also possible that Vps34 directly enhances endosomal cAMP production downstream of G_s activation (indicated by another red arrow with a question mark).

(Fig. 4C) This effect is specific to the isoproterenol-induced cAMP response because Vps34 inhibition did not attenuate the cAMP response caused by receptor-independent adenylyl cyclase activation with forskolin (Awad et al., 1983) (Fig. 4D). Thus, Vps34 activity not only supports long-term cellular β 2AR signaling, as indicated by its effect on resensitization, but it also plays an essential role in supporting the short-term signaling response.

Vps34 inhibition increases the endosomal β 2AR pool (Fig. 2C) and reduces the cAMP response after 10 minutes of isoproterenol application (Fig. 4C). One interpretation of these data is that the endosomal β 2AR pool cannot activate the cAMP response as strongly as the surface β 2AR pool. Indeed, our previous report suggested that endosomal β 2AR activation makes a relatively small, but functionally distinct, contribution to the cellular cAMP response (Irannejad et al., 2013; Tsvetanova and von Zastrow, 2014). It is also possible that the endosome signaling component is normally transient, as is the surface component, so that inhibiting recycling through Vps34 blockade causes the net response to dissipate. It will be interesting to investigate these possibilities in future studies.

The present results show that the endosomal system can indeed be targeted to manipulate canonical GPCR signaling, and identify an essential role of the endosomal PI3K Vps34 in promoting both short-term and long-term cellular responses mediated by a canonical GPCR cascade. Although class I PI3Ks are well known to promote signaling at the PM (Marat and Haucke, 2016), Vps34 has been studied primarily in the context of membrane trafficking, and its signaling functions are only beginning to emerge (Schink et al., 2013). Previous work on receptor-linked kinases, such as the transforming growth factor- β receptor, implicates Vps34 in recruiting PI3P-dependent signaling scaffolds to endosomes (Tsukazaki et al., 1998), and Vps34 was also shown to function as a noncanonical endosomal effector of yeast GPCR signaling (Slessareva et al., 2006). We propose that the present results, by establishing Vps34 as a druggable endosomal target that is essential for both short-term and sustained activation of canonical cellular GPCR signaling, open the door to the endosomal system as a target for future therapeutic intervention.

Acknowledgments

The authors thank all members of the von Zastrow laboratory for useful advice and critical discussion. The authors also thank A. Sundaram and D. Sheppard [University of California, San Francisco (UCSF)] for valuable discussion and providing reagents. A part of the live cell imaging experiments were performed in the Nikon Imaging Center at UCSF, directed by K. Thorn. The authors thank K. Thorn and D. Larson for essential advice and assistance. In addition, the authors thank S. Elmes in the Laboratory for Cell Analysis Core Facility at UCSF for technical assistance in the use of flow cytometry instruments.

Authorship Contributions

Participated in research design: Uchida, von Zastrow

Conducted experiments: Uchida

Contributed new reagents or analytic tools: Rataganira, Jullié, Shokat

Performed data analysis: Uchida, von Zastrow

Wrote or contributed to the writing of the manuscript: Uchida, Rutaganira, Jullié, Shokat, von Zastrow

References

Awad JA, Johnson RA, Jakobs KH, and Schultz G (1983) Interactions of forskolin and adenylate cyclase. Effects on substrate kinetics and protection against inactivation by heat and N-ethylmaleimide. *J Biol Chem* **258**:2960–2965.

Awad HO, Iyer V, Rosenfeld JL, Millman EE, Foster E, Moore RH, and Knoll BJ (2007) Inhibitors of phosphoinositide 3-kinase cause defects in the postendocytic sorting of beta2-adrenergic receptors. *Exp Cell Res* **313**: 2586–2596.

Bago R, Malik N, Munson MJ, Prescott AR, Davies P, Sommer E, Shpiro N, Ward R, Cross D, Ganley IG, et al. (2014) Characterization of VPS34-IN1, a selective inhibitor of Vps34, reveals that the phosphatidylinositol 3-phosphate-binding SGK3 protein kinase is a downstream target of class III phosphoinositide 3-kinase. *Biochem J* **463**:413–427.

Cao TT, Deacon HW, Reczek D, Bretscher A, and von Zastrow M (1999) A kinase-regulated PDZ-domain interaction controls endocytic sorting of the beta2-adrenergic receptor. *Nature* **401**:286–290.

Cullen PJ and Korswagen HC (2011) Sorting nexins provide diversity for retromer-dependent trafficking events. *Nat Cell Biol* **14**:29–37.

Dowdle WE, Nyfeler J, Nagel J, Elling RA, Liu S, Triantafellow E, Menon S, Wang Z, Honda A, Pardee G, et al. (2014) Selective VPS34 inhibitor blocks autophagy and uncovers a role for NCOA4 in ferritin degradation and iron homeostasis in vivo. *Nat Cell Biol* **16**:1069–1079.

Fan F, Binkowski BF, Butler BL, Stecha PF, Lewis MK, and Wood KV (2008) Novel genetically encoded biosensors using firefly luciferase. *ACS Chem Biol* **3**:346–351.

Fili N, Calleja V, Woscholski R, Parker PJ, and Larijani B (2006) Compartmental signal modulation: endosomal phosphatidylinositol 3-phosphate controls endosome morphology and selective cargo sorting. *Proc Natl Acad Sci USA* **103**:15473–15478.

Folkes AJ, Ahmadi K, Alderton WK, Alix S, Baker SJ, Box G, Chuckowree IS, Clarke PA, Depledge P, Eccles SA, et al. (2008) The identification of 2-(1H-indazol-4-yl)-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-4-morpholin-4-yl-thieno[3,2-d]pyrimidine (GDC-0941) as a potent, selective, orally bioavailable inhibitor of class I PI3 kinase for the treatment of cancer. *J Med Chem* **51**: 5522–5532.

Gillooly DJ, Morrow IC, Lindsay M, Gould R, Bryant NJ, Gaullier JM, Parton RG, and Stenmark H (2000) Localization of phosphatidylinositol 3-phosphate in yeast and mammalian cells. *EMBO J* **19**:4577–4588.

Goodman OB, Jr, Krupnick JG, Santini F, Gurevich VV, Penn RB, Gagnon AW, Keen JH, and Benovic JL (1996) Beta-arrestin acts as a clathrin adaptor in endocytosis of the beta2-adrenergic receptor. *Nature* **383**:447–450.

Hammond GR, Machner MP, and Balla T (2014) A novel probe for phosphatidylinositol 4-phosphate reveals multiple pools beyond the Golgi. *J Cell Biol* **205**: 113–126.

Hanyaloglu AC and von Zastrow M (2008) Regulation of GPCRs by endocytic membrane trafficking and its potential implications. *Annu Rev Pharmacol Toxicol* **48**: 537–568.

Herman PK and Emr SD (1990) Characterization of VPS34, a gene required for vacuolar protein sorting and vacuole segregation in *Saccharomyces cerevisiae*. *Mol Cell Biol* **10**:6742–6754.

Honda A, Harrington E, Cornella-Taracido I, Furet P, Knapp MS, Glick M, Triantafellow E, Dowdle WE, Wiedersheim D, Maniara W, et al. (2015) Potent, selective, and orally bioavailable inhibitors of VPS34 provide chemical tools to modulate autophagy in vivo. *ACS Med Chem Lett* **7**:72–76.

Irannejad R, Tomshine JC, Tomshine JR, Chevalier M, Mahoney JP, Steyaert J, Rasmussen SG, Sunahara RK, El-Samad H, Huang B, et al. (2013) Conformational biosensors reveal GPCR signalling from endosomes. *Nature* **495**:534–538.

Irannejad R, Tsvetanova NG, Lobingier BT, and von Zastrow M (2015) Effects of endocytosis on receptor-mediated signaling. *Curr Opin Cell Biol* **35**:137–143.

Irannejad R and von Zastrow M (2014) GPCR signaling along the endocytic pathway. *Curr Opin Cell Biol* **27**:109–116.

Jefferies HB, Cooke FT, Jat P, Boucheron C, Koizumi T, Hayakawa M, Kaizawa H, Ohishi T, Workman P, Waterfield MD, et al. (2008) A selective PIKfyve inhibitor blocks PtdIns(3,5)P2 production and disrupts endomembrane transport and retroviral budding. *EMBO Rep* **9**:164–170.

Knight ZA (2010) Small molecule inhibitors of the PI3-kinase family. *Curr Top Microbiol Immunol* **347**:263–278.

Laufer BE, Melerio C, Temkin P, Lei C, Hong W, Kortemme T, and von Zastrow M (2010) SNX27 mediates PDZ-directed sorting from endosomes to the plasma membrane. *J Cell Biol* **190**:565–574.

Lefkowitz RJ (2007) Seven transmembrane receptors: something old, something new. *Acta Physiol (Oxf)* **190**:9–19.

Marat AL and Haucke V (2016) Phosphatidylinositol 3-phosphates-at the interface between cell signalling and membrane traffic. *EMBO J* **35**:561–579.

McCartney AJ, Zhang Y, and Weisman LS (2014) Phosphatidylinositol 3,5-bisphosphate: low abundance, high significance. *BioEssays* **36**:52–64.

Morrison KJ, Moore RH, Carsrud ND, Trial J, Millman EE, Tuvim M, Clark RB, Barber R, Dickey BF, and Knoll BJ (1996) Repetitive endocytosis and recycling of the beta 2-adrenergic receptor during agonist-induced steady state redistribution. *Mol Pharmacol* **50**:692–699.

Murphy JE, Padilla BE, Hasdemir B, Cottrell GS, and Bunnett NW (2009) Endosomes: a legitimate platform for the signaling train. *Proc Natl Acad Sci USA* **106**: 17615–17622.

Naga Prasad SV, Barak LS, Rapacciuolo A, Caron MG, and Rockman HA (2001) Agonist-dependent recruitment of phosphoinositide 3-kinase to the membrane by beta-adrenergic receptor kinase 1. A role in receptor sequestration. *J Biol Chem* **276**:18953–18959.

Odley A, Hahn HS, Lynch RA, Marreez Y, Osinska H, Robbins J, and Dorn GW, 2nd (2004) Regulation of cardiac contractility by Rab4-modulated beta2-adrenergic receptor recycling. *Proc Natl Acad Sci USA* **101**:7082–7087.

Pierce KL, Premont RT, and Lefkowitz RJ (2002) Seven-transmembrane receptors. *Nat Rev Mol Cell Biol* **3**:639–650.

Pippig S, Andexinger S, and Lohse MJ (1995) Sequestration and recycling of beta 2-adrenergic receptors permit receptor resensitization. *Mol Pharmacol* **47**: 666–676.

Raiborg C, Schink KO, and Stenmark H (2013) Class III phosphatidylinositol 3-kinase and its catalytic product PtdIns3P in regulation of endocytic membrane traffic. *FEBS J* **280**:2730–2742.

Rasmussen SG, DeVree BT, Zou Y, Kruse AC, Chung KY, Kobilka TS, Thian FS, Chae PS, Pardon E, Calinski D, et al. (2011) Crystal structure of the β 2 adrenergic receptor-Gs protein complex. *Nature* **477**:549–555.

Ronan B, Flamand O, Vescovi L, Dureuil C, Durand L, Fassy F, Bachelot MF, Lamberton A, Mathieu M, Bertrand T, et al. (2014) A highly potent and selective Vps34 inhibitor alters vesicle trafficking and autophagy. *Nat Chem Biol* **10**: 1013–1019.

- Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, et al. (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682.
- Schink KO, Raiborg C, and Stenmark H (2013) Phosphatidylinositol 3-phosphate, a lipid that regulates membrane dynamics, protein sorting and cell signalling. *BioEssays* **35**:900–912.
- Shin HW, Hayashi M, Christoforidis S, Lacas-Gervais S, Hoepfner S, Wenk MR, Modregger J, Uttenweiler-Joseph S, Wilm M, Nystuen A, et al. (2005) An enzymatic cascade of Rab5 effectors regulates phosphoinositide turnover in the endocytic pathway. *J Cell Biol* **170**:607–618.
- Slessareva JE, Roult SM, Temple B, Bankaitis VA, and Dohlman HG (2006) Activation of the phosphatidylinositol 3-kinase Vps34 by a G protein alpha subunit at the endosome. *Cell* **126**:191–203.
- Sorensen SD, Linseman DA, McEwen EL, Heacock AM, and Fisher SK (1999) Inhibition of beta(2)-adrenergic and muscarinic cholinergic receptor endocytosis after depletion of phosphatidylinositol bisphosphate. *J Pharmacol Exp Ther* **290**:603–610.
- Sorkin A and von Zastrow M (2009) Endocytosis and signalling: intertwining molecular networks. *Nat Rev Mol Cell Biol* **10**:609–622.
- Temkin P, Lauffer B, Jäger S, Cimermancic P, Krogan NJ, and von Zastrow M (2011) SNX27 mediates retromer tubule entry and endosome-to-plasma membrane trafficking of signalling receptors. *Nat Cell Biol* **13**:715–721.
- Tsao PI and von Zastrow M (2000) Type-specific sorting of G protein-coupled receptors after endocytosis. *J Biol Chem* **275**:11130–11140.
- Tsukazaki T, Chiang TA, Davison AF, Attisano L, and Wrana JL (1998) SARA, a FYVE domain protein that recruits Smad2 to the TGFbeta receptor. *Cell* **95**:779–791.
- Tsvetanova NG and von Zastrow M (2014) Spatial encoding of cyclic AMP signaling specificity by GPCR endocytosis. *Nat Chem Biol* **10**:1061–1065.
- von Zastrow M and Williams JT (2012) Modulating neuromodulation by receptor membrane traffic in the endocytic pathway. *Neuron* **76**:22–32.
- Yu SS, Lefkowitz RJ, and Hausdorff WP (1993) Beta-adrenergic receptor sequestration. A potential mechanism of receptor resensitization. *J Biol Chem* **268**:337–341.
- Yudowski GA, Puthenveedu MA, Henry AG, and von Zastrow M (2009) Cargo-mediated regulation of a rapid Rab4-dependent recycling pathway. *Mol Biol Cell* **20**:2774–2784.

Address correspondence to: Yasunori Uchida or Mark von Zastrow, Department of Psychiatry, and Cellular and Molecular Pharmacology, University of California, San Francisco, 600 16th Street, San Francisco, CA 94143-2140. E-mail: uchida@med.kobe-u.ac.jp or mark.vonzastrow@ucsf.edu.
